



Homework 5

1. The free-free absorption coefficient of a pure hydrogen plasma is given by

$$\kappa_{\nu}^{\text{ff}} = n_e n_p \sigma_0 \nu^{-3} T^{-1/2} \left(1 - e^{-\frac{h\nu}{kT}} \right)$$

where n_e and n_p are the electron and proton number densities, respectively (given in cm^{-3}), ν is the frequency in Hz and T the temperature in K. In these units the value of σ_0 is then $\sigma_0 = 3.7 \times 10^8$.

Calculate the Rosseland mean opacity for κ_{ν}^{ff} . Discuss its temperature and density dependence. Compare with electron scattering:

$$\kappa_{e\text{-sc}} = n_e \sigma_e$$

where $\sigma_e = 6.6 \times 10^{-25} \text{ cm}^2$. Plot the borderline in the $(\log n_e, \log T)$ plane, where $\kappa_{e\text{-sc}}$ and $\bar{\kappa}^{\text{ff}}$ are equal and indicate where one is larger than the other.

(4 points)



Homework 5

2. In an ionized hydrogen plasma the total absorption coefficient is a combination of Thomson scattering and true absorption. Write a small subroutine and calculate

$$\kappa_R = \langle \kappa_\nu^{\text{ff}} + \kappa_{\text{e-sc}} \rangle_{\text{Rosseland}}$$

- Test the program using the results of Exercise 1. Discuss the values of κ_R at the borderline where $\kappa_{\text{e-sc}}$ and $\bar{\kappa}^{\text{ff}}$ are equal.

(3 points)

3. Calculate temperature, pressure and particle density stratifications in a stellar atmosphere with $T_{\text{eff}} = 20000 \text{ K}$ and $g = 10^4 \text{ cm s}^{-2}$ and $g = 10^2 \text{ cm s}^{-2}$.

Use: $T^4 = \frac{3}{4} T_{\text{eff}}^4 (\tau_R + 2/3)$ and $\frac{dP}{d\tau_R} = g \frac{\rho}{\kappa_R}$

as the basic equations.



Homework 5

Write a subroutine for numerical solution but discuss the solution analytically first for the case $\kappa_{e-sc} \gg \kappa_{\nu}^{ff}$.

Use τ_R (or its logarithm) as the depth scale.

Discuss the solutions for different gravity g .

Give electron densities at $\tau_R = 0.1, 1.0, 10.0$.

Assume $P = nkT$ for the equation of state with $n = n_e + n_p$ (n = total particle density, n_e = electron density, n_p = proton density) and $n_e = n_p$. k is the Boltzmann constant ($k = 1.38 \times 10^{-16}$ erg deg $^{-1}$).

To relate the mass density ρ to particle densities use $m_p = 1.67 \times 10^{-24}$ g for the proton mass.

(6 points)

Please return homework on Wednesday, October 24

!!! This is a lot of work. Start early !!!!