

# THE STAR FORMATION NEWSLETTER

*An electronic publication dedicated to early stellar evolution and molecular clouds*

No. 223 — 22 July 2011

Editor: Bo Reipurth (reipurth@ifa.hawaii.edu)

## *Abstracts of recently accepted papers*

### **Thermal Structure of a Protoplanetary Disk around HD163296: A Study of Vertical Temperature Distribution by CO Emission Lines**

**Eiji Akiyama<sup>1</sup>, Munetake Momose<sup>1</sup>, Hiroyuki Hayashi<sup>1</sup> and Yoshimi Kitamura<sup>2</sup>**

<sup>1</sup> College of Science, Ibaraki University, Bunkyo 2-1-1, Mito, Ibaraki, 310-8512, Japan

<sup>2</sup> Institute of Space and Astronomical Science, Japan Aerospace Exploration Agency, 3-1-1, Yoshinodai, Sagami-hara, Kanagawa, 229-8510, Japan

E-mail contact: 09nd401yat mcs.ibaraki.ac.jp

This paper presents observations of a protoplanetary disk around Herbig Ae star HD163296 in  $^{12}\text{CO}$  ( $J=1-0$ ),  $^{12}\text{CO}$  ( $J=3-2$ ),  $^{13}\text{CO}$  ( $J=1-0$ ), and  $^{13}\text{CO}$  ( $J=3-2$ ) emission lines. Double-peaked emission profiles originating from the rotating circumstellar disk were detected in all the lines. The disk parameters were estimated from model calculation in which the radial distribution of temperature or surface density inside the disk has a power-law form. The surface density should be sufficiently high so that the disk is optically thick for all the CO lines, as discussed in previous studies based on interferometric observations. The temperature and outer radius of the disk were also confirmed to be consistent with the previous results. Taking advantage of difference in position of the photosphere among the CO lines, we revealed temperature distribution in vertical direction. The temperature of  $^{12}\text{CO}$  ( $J=3-2$ ) emitting region is about twice higher than that of any other CO emitting region; the former is  $58.5 \pm 9.5$  K while the latter is  $31 \pm 15$  K at 100 AU from the central star, suggesting that there are at least two distinct temperature regions. The best fit temperature for  $^{13}\text{CO}$  ( $J=1-0$ ) that should trace the deepest region of the disk is even lower, implying that there is also a different temperature region in deep inside of the disk. Such vertical temperature distribution in a disk was identified both in T Tauri and Herbig Ae stars (e.g., DM Tau, AB Aur, and HD31648), and this should be a common feature in protoplanetary disks.

Accepted by PASJ

### **Rapid inward migration of planets formed by gravitational instability**

**Clément Baruteau<sup>1,2</sup>, Farzana Meru<sup>3,4</sup> and Sijme-Jan Paardekooper<sup>1</sup>**

<sup>1</sup> DAMTP, University of Cambridge, Wilberforce Road, Cambridge CB3 0WA, UK

<sup>2</sup> Department of Astronomy and Astrophysics, University of California, Santa Cruz, CA 95064, USA

<sup>3</sup> School of Physics, University of Exeter, Stocker Road, Exeter, EX4 4QL

<sup>4</sup> Institut für Astronomie und Astrophysik, Universität Tübingen, Auf der Mor genstelle 10, 72076 Tübingen, Germany

E-mail contact: C.Baruteau at damtp.cam.ac.uk

The observation of massive exoplanets at large separation ( $\geq 10$  AU) from their host star, like in the HR 8799 system, challenges theories of planet formation. A possible formation mechanism involves the fragmentation of massive self-gravitating discs into clumps. While the conditions for fragmentation have been extensively studied, little is known of the subsequent evolution of these giant planet embryos, in particular their expected orbital migration. Assuming a single planet has formed by fragmentation, we investigate its interaction with the gravitoturbulent disc it is embedded in. Two-dimensional hydrodynamical simulations are used with a simple prescription for the disc cooling. A steady gravitoturbulent disc is first set up, after which simulations are restarted including a planet with a range of masses approximately equal to the clump's initial mass expected in fragmenting discs. Planets rapidly migrate inwards, despite

the stochastic kicks due to the turbulent density fluctuations. We show that the migration timescale is essentially that of type I migration, with the planets having no time to open a gap. In discs with aspect ratio  $\sim 0.1$  at their forming location, planets with a mass comparable to, or larger than Jupiter's can migrate in as short as  $10^4$  years, that is, about 10 orbits at 100 AU. Massive planets formed at large separation from their star by gravitational instability are thus unlikely to stay in place, and should rapidly migrate towards the inner parts of protoplanetary discs, regardless of the planet mass.

Accepted by Monthly Notices of the Royal Astronomical Society

<http://arxiv.org/abs/1106.0487>

## Will the starless cores in Chamaeleon I and III turn prestellar?

Arnaud Belloche<sup>1</sup>, Bérengère Parise<sup>1</sup>, Frédéric Schuller<sup>1</sup>, Philippe André<sup>2</sup>, Sylvain Bontemps<sup>3</sup> and Karl M. Menten<sup>1</sup>

<sup>1</sup> Max-Planck Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany

<sup>2</sup> Laboratoire AIM, CEA/DSM-CNRS-Université Paris Diderot, IRFU/Service d'Astrophysique, CEA Saclay, 91191 Gif-sur-Yvette, France

<sup>3</sup> Université de Bordeaux, Laboratoire d'Astrophysique de Bordeaux, CNRS/INSU, UMR 5804, BP 89, 33271 Floirac cedex, France

E-mail contact: belloche at mpifr-bonn.mpg.de

*Context.* The nearby Chamaeleon molecular cloud complex is a good laboratory for studying the process of low-mass star formation because it consists of three clouds with very different properties. Chamaeleon III does not show any sign of star formation, while star formation has been very active in Chamaeleon I and may already be finishing.

*Aims.* Our goal is to determine whether star formation can proceed in Cha III by searching for prestellar cores, and to compare the results to our recent survey of Cha I.

*Methods.* We used the Large APEX Bolometer Array (LABOCA) to map Cha III in dust continuum emission at  $870\ \mu\text{m}$ . The map is compared with a 2MASS extinction map and decomposed with a multiresolution algorithm. The extracted sources are analyzed by carefully taking into account the spatial filtering inherent in the data reduction process.

*Results.* Twenty-nine sources are extracted from the  $870\ \mu\text{m}$  map, all of them starless. The estimated 90% completeness limit is  $0.18\ M_{\odot}$ . The starless cores are found down to a visual extinction of 1.9 mag, in marked contrast with other molecular clouds, including Cha I. Apart from this difference, the Cha III starless cores share very similar properties with those found in Cha I. They are less dense than those detected in continuum emission in other clouds by a factor of a few. At most two sources ( $< 7\%$ ) have a mass larger than the critical Bonnor-Ebert mass, which suggests that the fraction of prestellar cores is very low, even lower than in Cha I ( $< 17\%$ ). Only the most massive sources are candidate prestellar cores, in agreement with the correlation found earlier in the Pipe nebula. The mass distribution of the 85 starless cores of Cha I and III that are not candidate prestellar cores is consistent with a single power law down to the 90% completeness limit, with an exponent close to the Salpeter value. A fraction of the starless cores detected with LABOCA in Cha I and III may still grow in mass and become gravitationally unstable. Based on predictions of numerical simulations of turbulent molecular clouds, we estimate that at most 50% and 20% of the starless cores of Cha I and III, respectively, may form stars.

*Conclusions.* The LABOCA survey reveals that Cha III, and Cha I to some extent too, is a prime target to study the *formation* of prestellar cores, and thus the onset of star formation. Obtaining observational constraints on the duration of the core-building phase prior to gravitational collapse will be necessary to make further progress.

Accepted by Astronomy and Astrophysics

<http://de.arxiv.org/abs/1106.5064>

## Molecular outflows and hot molecular cores in G24.78+0.08 at sub-arcsecond angular resolution

M. T. Beltrán<sup>1</sup>, R. Cesaroni<sup>1</sup>, Q. Zhang<sup>2</sup>, R. Galván-Madrid<sup>2,3,4</sup>, H. Beuther<sup>5</sup>, C. Fallscheer<sup>5,6</sup>, R. Neri<sup>7</sup>, and C. Codella<sup>1</sup>

<sup>1</sup> INAF-Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125 Firenze, Italy

<sup>2</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge MA 02138, USA

<sup>3</sup> Centro de Radioastronomía y Astrofísica, Universidad Nacional Autónoma de México, Morelia 58090, Mexico

<sup>4</sup> Academia Sinica Institute of Astronomy and Astrophysics, Taipei 106, Taiwan

<sup>5</sup> Max Planck Institute for Astronomy, Königstuhl 17, 69117 Heidelberg, Germany

<sup>6</sup> Department of Physics and Astronomy, University of Victoria, 3800 Finnerty Road, Victoria, BC V8P 5C2, Canada

<sup>7</sup> IRAM, 300 Rue de la Piscine, F-38406 Saint Martin d'Hères, France

E-mail contact: mbeltran *at* arcetri.astro.it

*Context.* This study is part of a large project to study the physics of accretion and molecular outflows towards a selected sample of high-mass star-forming regions that show evidence of infall and rotation from previous studies.

*Aims.* We wish to make a thorough study at high-angular resolution of the structure and kinematics of the HMCs and corresponding molecular outflows in the high-mass star-forming region G24.78+0.08.

*Methods.* We carried out SMA and IRAM PdBI observations at 1.3 and 1.4 mm, respectively, of dust and of typical high-density and molecular outflow tracers with resolutions of  $<1''$ . Complementary IRAM 30-m  $^{12}\text{CO}$  and  $^{13}\text{CO}$  observations were carried out to recover the short spacing information of the molecular outflows.

*Results.* The millimeter continuum emission towards cores G24 A1 and A2 has been resolved into three and two cores, respectively, and named A1, A1b, A1c, A2, and A2b. All these cores are aligned in a southeast-northwest direction coincident with that of the molecular outflows detected in the region, which suggests a preferential direction for star formation in this region. The masses of the cores range from 7 to 22  $M_{\odot}$ , and the rotational temperatures from 128 to 180 K. The high-density tracers have revealed the existence of two velocity components towards A1. One of them peaks close to the position of the millimeter continuum peak and of the HC HII region and is associated with the velocity gradient seen in  $\text{CH}_3\text{CN}$  towards this core, while the other one peaks southwest of core A1 and is not associated with any millimeter continuum emission peak. The position-velocity plots along outflow A and the  $^{13}\text{CO}$  (2–1) averaged blueshifted and redshifted emission indicate that this outflow is driven by core A2. Core A1 apparently does not drive any outflow. The knotty appearance of the highly collimated outflow C and the  $^{12}\text{CO}$  position-velocity plot suggest an episodic outflow, where the knots are made of swept-up ambient gas.

Accepted by A&A

<http://arxiv.org/abs/1107.0314>

## Photoelectric cross-sections in protoplanetary disks

Thomas J. Bethell<sup>1</sup> and Edwin A. Bergin<sup>1</sup>

<sup>1</sup> University of Michigan - Ann Arbor, USA

E-mail contact: tbethell *at* umich.edu

We provide simple polynomial fits to the X-ray photoelectric cross-sections ( $0.03 < E < 10\text{keV}$ ) for mixtures of gas and dust found in protoplanetary disks. Using the solar elemental abundances of Asplund et al. (2009) we treat the gas and dust components separately, facilitating the further exploration evolutionary processes such as grain settling and grain growth. We find that blanketing due to advanced grain-growth ( $a_{max} > 1\mu\text{m}$ ) can reduce the X-ray opacity of dust appreciably at  $E_X \sim 1\text{keV}$ , coincident with the peak of typical T Tauri X-ray spectra. However, the reduction of dust opacity by dust settling, which is known to occur in protoplanetary disks, is probably a more significant effect. The absorption of 1-10keV X-rays is dominated by gas opacity once the dust abundance has been reduced to about 1% of its diffuse interstellar value. The gas disk establishes a floor to the opacity at which point X-ray transport becomes insensitive to further dust evolution. Our choice of fitting function follows that of Morrison & McCammon (1983), providing a degree of backward-compatibility.

Accepted by ApJ

## Propagation of Ly- $\alpha$ in protoplanetary disks

Thomas J. Bethell<sup>1</sup> and Edwin A. Bergin<sup>1</sup>

<sup>1</sup> University of Michigan - Ann Arbor, USA

E-mail contact: tbethell *at* umich.edu

We study the role resonant scattering plays in the transport of Ly- $\alpha$  photons in accreting protoplanetary disk systems

subject to varying degrees of dust settling. While the intrinsic stellar FUV spectrum of accreting T Tauri systems may already be dominated by a strong, broad Ly- $\alpha$  line ( $\sim 80\%$  of the FUV luminosity), we find that resonant scattering further enhances the Ly- $\alpha$  density in the deep molecular layers of the disk. Ly- $\alpha$  is scattered downwards efficiently by the photodissociated atomic hydrogen layer that exists above the molecular disk. In contrast, FUV-continuum photons pass unimpeded through the photodissociation layer, and (forward-)scatter inefficiently off dust grains. Using detailed, adaptive grid Monte Carlo radiative transfer simulations we show that the resulting Ly- $\alpha$ /FUV-continuum photon density ratio is strongly stratified; FUV-continuum dominated in the photodissociation layer, and Ly- $\alpha$  dominated field in the molecular disk. The enhancement is greatest in the interior of the disk ( $r \sim 1\text{AU}$ ) but is also observed in the outer disk ( $r \sim 100\text{AU}$ ). The majority of the total disk mass is shown to be increasingly Ly- $\alpha$  dominated as dust settles towards the midplane.

Accepted by ApJ

## The Coalsack near and far

**H. Beuther<sup>1</sup>, J. Kainulainen<sup>1</sup>, Th. Henning<sup>1</sup>, R. Plume<sup>2</sup> and F. Heitsch<sup>3</sup>**

<sup>1</sup> Max-Planck-Institute for Astronomy, Königstuhl 17, 69117 Heidelberg, Germany

<sup>2</sup> Department of Physics and Astronomy, University of Calgary, Calgary, Canada

<sup>3</sup> Department of Physics & Astronomy, University of North Carolina at Chapel Hill, CB 3255, Chapel Hill, NC 27599-3255, USA

E-mail contact: beuther *at* mpia.de

*Context:* The large Coalsack dark cloud is one of the most prominent southern starless clouds, which is even visible to the naked eye. Furthermore, it is one of the rare molecular clouds without clear signs of star formation.

*Aims:* We investigate the dynamical properties of the gas within the Coalsack.

*Methods:* The two highest extinction regions were mapped with the APEX telescope in  $^{13}\text{CO}(2-1)$  comprising a region of  $\sim 1$  square degree.

*Results:* In addition to the well-known, nearby gas component around  $-4\text{ km s}^{-1}$ , we identified additional molecular gas components – in particular a second extended molecular cloud at a velocity of  $\sim -30\text{ km s}^{-1}$  and an estimated distance of  $\sim 3.1\text{ kpc}$  – that dominate the column density and visual extinction distributions in the northeastern part of the Coalsack. Although comprising  $\sim 2600\text{ M}_{\odot}$ , the mass of this distant cloud is distributed over an extent of  $\sim 73\text{ pc}$ , much larger than typical high-mass infrared dark clouds. Its filamentary structure is consistent with a compressible gaseous self-gravitating cylinder, and its low mass per length indicates that it may be stable against gravitational collapse. We find barely any mid-infrared emission in archival MSX data, which is indicative of almost no star-formation activity in the near and far cloud complexes. The nearby clouds have narrow, almost thermal velocity dispersions with median values between  $0.2$  and  $0.4\text{ km s}^{-1}$ , which is also consistent with low star-formation activity. Only Tapia’s Globule 2 exhibits a velocity dispersion increase toward the extinction peak and peak-velocity gradients over the core, which is indicative of a state of elevated dynamical properties.

*Conclusions:* The Coalsack is not one single coherent structure, but consists of several cloud complexes nearby as well as at several kpc distance. All studied clouds appear as starless low-turbulence regions that may not even collapse in the future. Only one globule exhibits more dynamical signatures and is a good candidate for present/future star formation.

Accepted by Astronomy & Astrophysics

<http://www.mpia-hd.mpg.de/homes/beuther/papers.html>

## Herschel observations of the Herbig-Haro objects HH52-54

**P. Bjerkeli<sup>1</sup>, R. Liseau<sup>1</sup>, B. Nisini<sup>2</sup>, M. Tafalla<sup>3</sup>, M. Benedettini<sup>4</sup>, P. Bergman<sup>1</sup>, O. Dionatos<sup>5</sup>, T. Giannini<sup>2</sup>,**

**G. Herczeg<sup>6</sup>, K. Justtanont<sup>1</sup>, B. Larsson<sup>7</sup>, C. M<sup>c</sup>Coey<sup>8</sup>, M. Olberg<sup>1</sup> and A.O.H Olofsson<sup>1</sup>**

<sup>1</sup> Department of Earth and Space Sciences, Chalmers University of Technology, Onsala Space Observatory, 439 92 Onsala, Sweden

<sup>2</sup> INAF - Osservatorio Astronomico di Roma, Via di Frascati 33, 00040 Monte Porzio Catone, Italy

<sup>3</sup> Observatorio Astronómico Nacional (IGN), Calle Alfonso XII,3. 28014, Madrid, Spain

<sup>4</sup> INAF - Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, 50125 Firenze, Italy

<sup>5</sup> Centre for Star and Planet Formation, Natural History Museum of Denmark, University of Copenhagen, Øster Voldgade 5-7, 1350 Copenhagen, Denmark

<sup>6</sup> Max Planck Institut for Extraterrestrische Physik, Garching, Germany

<sup>7</sup> Department of Astronomy, Stockholm University, AlbaNova, 106 91 Stockholm, Sweden

<sup>8</sup> University of Waterloo, Department of Physics and Astronomy, Waterloo, Ontario, Canada

E-mail contact: per.bjerkeli *at* chalmers.se

*Context.* The emission from Herbig-Haro objects and supersonic molecular outflows is understood as cooling radiation behind shocks, initiated by a (proto-)stellar wind or jet. Within a given object, one often observes the occurrence of both dissociative (J-type) and non-dissociative (C-type) shocks, owing to the collective effects of internally varying shock velocities.

*Aims.* We are aiming at the observational estimation of the relative contribution to the cooling by CO and H<sub>2</sub>O, as this provides decisive information for the understanding of the oxygen chemistry behind interstellar shock waves.

*Methods.* The high sensitivity of HIFI, in combination with its high spectral resolution capability, allows us to trace the H<sub>2</sub>O outflow wings at unprecedented signal-to-noise. From the observation of spectrally resolved H<sub>2</sub>O and CO lines in the HH52-54 system, both from space and from ground, we arrive at the spatial and velocity distribution of the molecular outflow gas. Solving the statistical equilibrium and non-LTE radiative transfer equations provides us with estimates of the physical parameters of this gas, including the cooling rate ratios of the species. The radiative transfer is based on an Accelerated Lambda Iteration code, where we use the fact that variable shock strengths, distributed along the front, are naturally implied by a curved surface.

*Results.* Based on observations of CO and H<sub>2</sub>O spectral lines, we conclude that the emission is confined to the HH54 region. The quantitative analysis of our observations favours a ratio of the CO-to-H<sub>2</sub>O-cooling-rate  $\gg 1$ . Formally, we derive the ratio  $\Lambda(\text{CO})/\Lambda(o\text{-H}_2\text{O}) = 10$ , which is in good agreement with earlier determination of 7 based on ISO-LWS observations. From the best-fit model to the CO emission, we arrive at an H<sub>2</sub>O abundance close to  $1 \times 10^{-5}$ . The line profiles exhibit two components, one of which is triangular and another, which is a superposed, additional feature. This additional feature likely originates from a region smaller than the beam where the *ortho*-water abundance is smaller than in the quiescent gas.

*Conclusions.* Comparison with recent shock models indicate that a planar shock can not easily explain the observed line strengths and triangular line profiles. We conclude that the geometry can play an important role. Although abundances support a scenario where J-type shocks are present, higher cooling rate ratios than predicted by these type of shocks are derived.

Accepted by Astronomy & Astrophysics

<http://arxiv.org/abs/1106.5440>

## Evolution of the Solar Nebula. IX. Gradients in the Spatial Heterogeneity of the Short-Lived Radioisotopes <sup>60</sup>Fe and <sup>26</sup>Al and the Stable Oxygen Isotopes

Alan P. Boss<sup>1</sup>

<sup>1</sup> DTM, Carnegie Institution, USA

E-mail contact: boss *at* dtm.ciw.edu

Short-lived radioisotopes (SLRI) such as <sup>60</sup>Fe and <sup>26</sup>Al were likely injected into the solar nebula in a spatially and temporally heterogeneous manner. Marginally gravitationally unstable (MGU) disks, of the type required to form gas giant planets, are capable of rapid homogenization of isotopic heterogeneity as well as of rapid radial transport of dust grains and gases throughout a protoplanetary disk. Two different types of new models of a MGU disk in orbit around a solar-mass protostar are presented. The first set has variations in the number of terms in the spherical harmonic solution for the gravitational potential, effectively studying the effect of varying the spatial resolution of the gravitational torques responsible for MGU disk evolution. The second set explores the effects of varying the initial minimum value of the Toomre  $Q$  stability parameter, from values of 1.4 to 2.5, i.e., toward increasingly less unstable disks. The new models show that the basic results are largely independent of both sets of variations. MGU disk models robustly result in rapid mixing of initially highly heterogeneous distributions of SLRIs to levels of  $\sim 10\%$  in both the inner ( $< 5$  AU) and outer ( $> 10$  AU) disk regions, and to even lower levels ( $\sim 2\%$ ) in intermediate regions, where gravitational torques are most effective at mixing. These gradients should have cosmochemical implications

for the distribution of SLRIs and stable oxygen isotopes contained in planetesimals (e.g., comets) formed in the giant planet region ( $\sim 5$  to  $\sim 10$  AU) compared to those formed elsewhere.

Accepted by ApJ

<http://www.dtm.ciw.edu/users/boss/ftp/nebulaix.pdf>

## A survey for near-infrared H<sub>2</sub> emission in Herbig Ae/Be stars: emission from the outer disks of HD 97048 and HD 100546

A. Carmona<sup>1,2</sup>, G. van der Plas<sup>3</sup>, M.E. van den Ancker<sup>4</sup>, M. Audard<sup>1,2</sup>, L.B.F.M Waters<sup>5,3</sup>, D. Fedele<sup>6</sup>, B. Acke<sup>7</sup> and E. Pantin<sup>8</sup>

<sup>1</sup> ISDC Data Centre for Astrophysics, University of Geneva, chemin d'Ecogia 16, 1290 Versoix, Switzerland

<sup>2</sup> Observatoire de Genève, University of Geneva, chemin des Maillettes 51, 1290 Sauverny, Switzerland

<sup>3</sup> Sterrenkundig Instituut 'Anton Pannekoek', University of Amsterdam, Kruislaan 403, 1098 SJ Amsterdam, The Netherlands

<sup>4</sup> European Southern Observatory, Karl-Schwarzschild-Str.2, D 85748 Garching bei München, Germany

<sup>5</sup> SRON Netherlands Institute for Space Research, Sorbonnelaan 2, 3584 CA Utrecht, The Netherlands

<sup>6</sup> Department of Physics and Astronomy, Johns Hopkins University, Baltimore, MD 21218, USA

<sup>7</sup> Instituut voor Sterrenkunde, K. U. Leuven, Celestijnenlaan 200D, 3001 Leuven, Belgium

<sup>8</sup> Service d'Astrophysique, CEA Saclay, F-91191 Gif-sur-Yvette, Cedex, France

E-mail contact: andres.carmona *at* unige.ch

We report on a sensitive search for H<sub>2</sub> 1-0 S(1), 1-0 S(0) and 2-1 S(1) ro-vibrational emission at 2.12, 2.22 and 2.25  $\mu\text{m}$  in a sample of 15 Herbig Ae/Be stars employing CRIRES, the ESO-VLT near-infrared high-resolution spectrograph, at R $\sim$ 90,000. We report the detection of the H<sub>2</sub> 1-0 S(1) line toward HD 100546 and HD 97048. In the other 13 targets, the line is not detected. The H<sub>2</sub> 1-0 S(0) and 2-1 S(1) lines are undetected in all sources. These observations are the first detection of near-IR H<sub>2</sub> emission in HD 100546. The H<sub>2</sub> 1-0 S(1) lines observed in HD 100546 and HD 97048 are observed at a velocity consistent with the rest velocity of both stars, suggesting that they are produced in the circumstellar disk. In HD 97048, the emission is spatially resolved and it is observed to extend at least up to 200 AU from the star. We report an increase of one order of magnitude in the H<sub>2</sub> 1-0 S(1) line flux with respect to previous measurements taken in 2003 for this star, which suggests line variability. In HD 100546 the emission is tentatively spatially resolved and may extend at least up to 50 AU from the star. Modeling of the H<sub>2</sub> 1-0 S(1) line profiles and their spatial extent with flat keplerian disks shows that most of the emission is produced at a radius larger than 5 AU. Upper limits to the H<sub>2</sub> 1-0 S(0)/ 1-0 S(1) and H<sub>2</sub> 2-1 S(1)/1-0 S(1) line ratios in HD 97048 are consistent with H<sub>2</sub> gas at T>2000 K and suggest that the emission observed may be produced by X-ray excitation. The upper limits for the line ratios for HD 100546 are inconclusive. Because the H<sub>2</sub> emission is located at large radii, for both sources a thermal emission scenario (i.e., gas heated by collisions with dust) is implausible. We argue that the observation of H<sub>2</sub> emission at large radii may be indicative of an extended disk atmosphere at radii > 5 AU. This may be explained by a hydrostatic disk in which gas and dust are thermally decoupled or by a disk wind caused by photoevaporation.

Accepted by A&A.

<http://arxiv.org/abs/1107.1994>

## Dissecting a hot molecular core: The case of G31.41+0.31

R. Cesaroni<sup>1</sup>, M.T. Beltrán<sup>1</sup>, Q. Zhang<sup>2</sup>, H. Beuther<sup>3</sup>, and C. Fallscheer<sup>3,4</sup>

<sup>1</sup> Osservatorio Astrofisico di Arcetri, INAF, Largo E. Fermi 5, I-50125 Firenze, Italy

<sup>2</sup> Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138, U.S.A.

<sup>3</sup> Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany

<sup>4</sup> Department of Physics and Astronomy, University of Victoria, 3800 Finnerty Road, Victoria, BC V8P 5C2, Canada

E-mail contact: cesa *at* arcetri.astro.it

*Context* The role of disks in the formation of high-mass stars is still a matter of debate but the detection of circumstellar disks around O-type stars would have a profound impact on high-mass star formation theories.

*Aims* We made a detailed observational analysis of a well known hot molecular core lying in the high-mass star-forming

region G31.41+0.31. This core is believed to contain deeply embedded massive stars and presents a velocity gradient that has been interpreted either as rotation or as expansion, depending on the authors. Our aim was to shed light on this question and possibly prepare the ground for higher resolution ALMA observations which could directly detect circumstellar disks around the embedded massive stars.

*Methods* Observations at sub-arcsecond resolution were performed with the Submillimeter Array in methyl cyanide, a typical hot molecular core tracer, and  $^{12}\text{CO}$  and  $^{13}\text{CO}$ , well known outflow tracers. We also obtained sensitive continuum maps at 1.3 mm.

*Results* Our findings confirm the existence of a sharp velocity gradient across the core, but cannot confirm the existence of a bipolar outflow perpendicular to it. The improved angular resolution and sampling of the uv plane allow us to attain higher quality channel maps of the  $\text{CH}_3\text{CN}$  lines with respect to previous studies and thus significantly improve our knowledge of the structure and kinematics of the hot molecular core.

*Conclusions* While no conclusive argument can rule out any of the two interpretations (rotation or expansion) proposed to explain the velocity gradient observed in the core, in our opinion the observational evidence collected so far indicates the rotating toroid as the most likely scenario. The outflow hypothesis appears less plausible, because the dynamical time scale is too short compared to that needed to form species such as  $\text{CH}_3\text{CN}$ , and the mass loss and momentum rates estimated from our measurements appear too high.

Accepted by Astron. Astrophys.

[http://www.arcetri.astro.it/~starform/preprints/cesa\\_22.pdf](http://www.arcetri.astro.it/~starform/preprints/cesa_22.pdf)

## Radio Imaging of the NGC 1333 IRAS 4A Region: Envelope, Disks, and Outflows of a Protostellar Binary System

Minho Choi<sup>1</sup>, Miju Kang<sup>1</sup>, Ken'ichi Tatematsu<sup>2</sup>, Jeong-Eun Lee<sup>3</sup> and Geumsook Park<sup>4</sup>

<sup>1</sup> Korea Astronomy and Space Science Institute, 776 Daedeokdaero, Yuseong, Daejeon 305-348, South Korea

<sup>2</sup> National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

<sup>3</sup> Department of Astronomy and Space Science, Kyung Hee University, Yongin, Gyeonggi 446-701, South Korea

<sup>4</sup> Department of Physics and Astronomy, Seoul National University, Seoul 151-742, South Korea

E-mail contact: minho *at* kasi.re.kr

The NGC 1333 IRAS 4A protobinary was observed in the 1.3 cm and 6.9 mm continuum and the ammonia and SiO lines, with an angular resolution of about 0.4 arcseconds. The continuum maps show the circumstellar structures of the two protostars, A1 and A2. The A1 system is brighter and more massive than the A2 system. The ratio of mass, including dense gas and protostar, is about 6. The properties of the circumstellar disks and outflows suggest that A1 may be younger than A2. The deflected part of the northeastern jet of A2 is bright in the SiO line, and the distance between the brightest peak and deflection point suggests that the enhancement of SiO takes about 100 yr after the collision with a dense core. The ammonia maps show a small structure that seems to be a part of the obstructing core. The outflow properties were studied by comparing interferometric maps of SiO, ammonia, formaldehyde, and HCN lines. Their overall structures agree well, suggesting that these species are excited by the same mechanism. However, the intensity distributions show that SiO is chemically unique. SiO may be directly linked to the primary jet while the other species may be tracing the entrained ambient gas.

Accepted by PASJ

<http://arxiv.org/abs/1107.3877>

## Clues to the star formation in NGC 346 across time and space

Guido De Marchi<sup>1</sup>, Nino Panagia<sup>2,3,4</sup> and Elena Sabbi<sup>2</sup>

<sup>1</sup> European Space Agency, Space Science Department, Keplerlaan 1, 2200 AG Noordwijk, Netherlands

<sup>2</sup> Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

<sup>3</sup> INAF-CT, Osservatorio Astrofisico di Catania, Via S. Sofia, 78, 95123 Catania, Italy

<sup>4</sup> Supernova Limited, OYV #131, Northsound Rd., Virgin Gorda, British Virgin Islands

E-mail contact: gdemarchi *at* rssd.esa.int

We have studied the properties of the stellar populations in the field of the NGC 346 cluster in the Small Magellanic

Cloud, using the results of a novel self-consistent method that provides a reliable identification of pre-main sequence (PMS) objects actively undergoing mass accretion, regardless of their age. The 680 identified bona-fide PMS stars show a bimodal age distribution, with two roughly equally numerous populations peaked respectively at  $\sim 1$  Myr, and  $\sim 20$  Myr. We use the age and other physical properties of these PMS stars to study how star formation has proceeded across time and space in NGC 346. We find no correlation between the locations of young and old PMS stars, nor do we find a correspondence between the positions of young PMS stars and those of massive OB stars of similar age. Furthermore, the mass distribution of stars with similar age shows large variations throughout the region. We conclude that, while on a global scale it makes sense to talk about an initial mass function, this concept is not meaningful for individual star-forming regions. An interesting implication of the separation between regions where massive stars and low-mass objects appear to form is that high-mass stars might not be “perfect” indicators of star formation and hence a large number of low-mass stars formed elsewhere might have so far remained unnoticed. For certain low surface density galaxies this way of preferential low-mass star formation may be the predominant mechanism, with the consequence that their total mass as derived from the luminosity may be severely underestimated and that their evolution is not correctly understood.

Accepted by The Astrophysical Journal

<http://arxiv.org/abs/1106.5780>

## Photometric determination of the mass accretion rates of pre-main sequence stars. II. NGC346 in the Small Magellanic Cloud

Guido De Marchi<sup>1</sup>, Nino Panagia<sup>2,3,4</sup>, Martino Romaniello<sup>5</sup>, Elena Sabbi<sup>2</sup>, Marco Sirianni<sup>1</sup>, Pier Giorgio Prada Moroni<sup>6,7</sup> and Scilla Degl’Innocenti<sup>6,7</sup>

<sup>1</sup> European Space Agency, Space Science Department, Keplerlaan 1, Noordwijk, Netherlands

<sup>2</sup> Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

<sup>3</sup> INAF–CT, Osservatorio Astrofisico di Catania, Via S. Sofia 78, 95123 Catania, Italy

<sup>4</sup> Supernova Limited, OYV # 131, Northsound Rd., Virgin Gorda, British Virgin Islands

<sup>5</sup> European Southern Observatory, Karl Schwarzschild-Str. 2, 85748 Garching, Germany

<sup>6</sup> Dipartimento di Fisica Enrico Fermi, Università di Pisa, Largo Pontecorvo 3, 56127 Pisa, Italy

<sup>7</sup> INFN, Sezione di Pisa, Largo Pontecorvo 3, 56127 Pisa, Italy

E-mail contact: [gdemarchi@rssd.esa.int](mailto:gdemarchi@rssd.esa.int)

We have studied the properties of the stellar populations in the field of the NGC 346 cluster in the Small Magellanic Cloud, using a novel self-consistent method that allows us to reliably identify pre-main sequence (PMS) objects actively undergoing mass accretion, regardless of their age. The method does not require spectroscopy and combines broad-band  $V$  and  $I$  photometry with narrow-band  $H\alpha$  imaging to identify all stars with excess  $H\alpha$  emission and derive the accretion luminosity  $L_{\text{acc}}$  and mass accretion rate  $\dot{M}_{\text{acc}}$  for all of them. The application of this method to existing HST/ACS photometry of the NGC 346 field has allowed us to identify and study 680 bona-fide PMS stars with masses from  $\sim 0.4 M_{\odot}$  to  $\sim 4 M_{\odot}$  and ages in the range from  $\sim 1$  Myr to  $\sim 30$  Myr. Previous investigations of this region, based on the same data, had identified young ( $\sim 3$  Myr old) candidate PMS stars on the basis of their broad-band colours. In this study we show that there are at least two, almost equally numerous, young populations with distinct ages of respectively  $\sim 1$  and  $\sim 20$  Myr. We provide for all of them accurate physical parameters.

We take advantage of the unprecedented size of our PMS sample and of its spread in mass and age to study the evolution of the mass accretion rate as a function of stellar parameters. We find that, regardless of stellar mass, the mass accretion rate decreases with roughly the square root of the age, or about three times slower than predicted by current models of viscous disc evolution, and that more massive stars have systematically higher mass accretion rate in proportion to their mass. A multivariate linear regression fit reveals that  $\log \dot{M}_{\text{acc}} \simeq -0.6 \log t + \log m + c$ , where  $t$  is the age of the star,  $m$  its mass and  $c$  a quantity that is higher at lower metallicity. This result is consistent with measurements of the mass accretion rate in the 30 Dor region and in the Milky Way and suggests that longer duration for mass accretion could be related to lower metallicity. The high mass accretion rates that we find suggest that a considerable amount of mass is accreted during the PMS phase, of order  $\sim 0.2 M_{\odot}$  or possibly  $\sim 20\%$  of the final mass for stars with mass  $m < 1 M_{\odot}$  if their discs are eroded by 20 Myr, i.e. before they reach the main sequence. Therefore, PMS evolutionary models that do not account for this effect will systematically underestimate the true age when compared with the observations.



## Star formation in 30 Doradus

Guido De Marchi<sup>1</sup>, Francesco Paresce<sup>2</sup>, Nino Panagia<sup>3,4,5</sup>, Giacomo Beccari<sup>6</sup>, Loredana Spezzi<sup>1</sup>, Marco Sirianni<sup>1</sup>, Morten Andersen<sup>1</sup>, Max Mutchler<sup>3</sup>, Bruce Balick<sup>7</sup>, Michael A. Dopita<sup>8,9,10</sup>, Jay A. Frogel<sup>11</sup>, Bradley C. Whitmore<sup>3</sup>, Howard Bond<sup>3</sup>, Daniela Calzetti<sup>12</sup>, C. Marcella Carollo<sup>13</sup>, Michael J. Disney<sup>14</sup>, Donald N. B. Hall<sup>15</sup>, Jon A. Holtzman<sup>16</sup>, Randy A. Kimble<sup>17</sup>, Patrick J. McCarthy<sup>18</sup>, Robert W. O’Connell<sup>19</sup>, Abhijit Saha<sup>20</sup>, Joseph I. Silk<sup>21</sup>, John T. Trauger<sup>22</sup>, Alistair R. Walker<sup>23</sup>, Rogier A. Windhorst<sup>24</sup>, and Erick T. Young<sup>25</sup>

<sup>1</sup> European Space Agency, Space Science Department, Keplerlaan 1, 2200 AG Noordwijk, Netherlands

<sup>2</sup> Istituto di Astrofisica Spaziale e Fisica Cosmica, Via Gobetti 101, 40129 Bologna, Italy

<sup>3</sup> Space Telescope Science Institute, 3700 San Martin Drive, Baltimore MD 21218, USA

<sup>4</sup> INAF–CT, Osservatorio Astrofisico di Catania, Via S. Sofia 78, 95123 Catania, Italy

<sup>5</sup> Supernova Limited, OYV #131, Northsound Rd., Virgin Gorda, British Virgin Islands

<sup>6</sup> European Southern Observatory, Karl–Schwarzschild–Str. 2, 85748 Garching, Germany

<sup>7</sup> Department of Astronomy, University of Washington, Seattle, WA 98195–1580, USA

<sup>8</sup> Mount Stromlo and Siding Spring Observatories, Research School of Astronomy and Astrophysics, Australian National University, Cotter Road, Weston Creek, ACT 2611, Australia

<sup>9</sup> Astronomy Department, King Abdulaziz University, P.O. Box 80203, Jeddah, Saudi Arabia

<sup>10</sup> Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822

<sup>11</sup> Galaxies Unlimited, 8726 Hickory Bend Trail, Potomac, MD 20854, USA

<sup>12</sup> Department of Astronomy, University of Massachusetts, Amherst, MA 01003, USA

<sup>13</sup> Department of Physics, ETH–Zurich, Zurich, 8093, Switzerland

<sup>14</sup> School of Physics and Astronomy, Cardiff University, Cardiff CF24 3AA, United Kingdom

<sup>15</sup> Institute for Astronomy, University of Hawaii, Honolulu, HI 96822, USA

<sup>16</sup> Department of Astronomy, New Mexico State University, Las Cruces, NM 88003, USA

<sup>17</sup> NASA–Goddard Space Flight Center, Greenbelt, MD 20771, USA

<sup>18</sup> Observatories of the Carnegie Institution of Washington, Pasadena, CA 91101–1292, USA

<sup>19</sup> Department of Astronomy, University of Virginia, Charlottesville, VA 22904–4325, USA

<sup>20</sup> National Optical Astronomy Observatories, Tucson, AZ 85726–6732, USA

<sup>21</sup> Department of Physics, University of Oxford, Oxford OX1 3PU, United Kingdom

<sup>22</sup> NASA–Jet Propulsion Laboratory, Pasadena, CA 91109, USA

<sup>23</sup> Cerro Tololo Inter–American Observatory, La Serena, Chile

<sup>24</sup> School of Earth and Space Exploration, Arizona State University, Tempe, AZ 85287–1404, USA

<sup>25</sup> SOFIA Science Center, NASA Ames Research Center, Moffett Field, California 94035, USA

E-mail contact: [gdemarchi@rssd.esa.int](mailto:gdemarchi@rssd.esa.int)

Using observations obtained with the Wide Field Camera 3 (WFC3) on board the *Hubble Space Telescope* (HST), we have studied the properties of the stellar populations in the central regions of 30 Dor, in the Large Magellanic Cloud. The observations clearly reveal the presence of considerable differential extinction across the field. We characterise and quantify this effect using young massive main sequence stars to derive a statistical reddening correction for most objects in the field. We then search for pre-main sequence (PMS) stars by looking for objects with a strong ( $> 4\sigma$ )  $H\alpha$  excess emission and find about 1150 of them over the entire field. Comparison of their location in the Hertzsprung–Russell diagram with theoretical PMS evolutionary tracks for the appropriate metallicity reveals that about one third of these objects are younger than  $\sim 4$  Myr, compatible with the age of the massive stars in the central ionising cluster R 136, whereas the rest have ages up to  $\sim 30$  Myr, with a median of  $\sim 12$  Myr. This indicates that star formation has proceeded over an extended period of time, although we cannot discriminate between an extended episode and a series of short and frequent bursts that are not resolved in time. While the younger PMS population preferentially occupies the central regions of the cluster, older PMS objects are more uniformly distributed across the field and are remarkably few at the very centre of the cluster. We attribute this latter effect to photoevaporation of the older circumstellar discs caused by the massive ionising members of R 136.

## Feedback Regulated Star Formation: Implications for the Kennicutt-Schmidt Law

Sami Dib<sup>1</sup>

<sup>1</sup> Imperial College London, Blackett Laboratory, Prince Consort Road, SW7 2AZ, London, United Kingdom

E-mail contact: [s.dib@imperial.ac.uk](mailto:s.dib@imperial.ac.uk)

We derive a metallicity dependent relation between the surface density of the star formation rate ( $\Sigma_{SFR}$ ) and the gas surface density ( $\Sigma_g$ ) in a feedback regulated model of star formation in galactic disks. In this model, star formation occurs in gravitationally bound protocluster clumps embedded in larger giant molecular clouds with the protocluster clump mass function following a power law function with a slope of  $-2$ . Metallicity dependent feedback is generated by the winds of OB stars ( $M \gtrsim 5 M_\odot$ ) that form in the clumps. The quenching of star formation in clumps of decreasing metallicity occurs at later epochs due to weaker wind luminosities, thus resulting in higher final star formation efficiencies ( $SFE_{exp}$ ). By combining  $SFE_{exp}$  with the timescales on which gas expulsion occurs, we derive the metallicity dependent star formation rate per unit time in this model as a function of  $\Sigma_g$ . This is combined with the molecular gas fraction in order to derive the global dependence of  $\Sigma_{SFR}$  on  $\Sigma_g$ . The model reproduces very well the observed star formation laws extending from low gas surface densities up to the starburst regime. Furthermore, our results show a dependence of  $\Sigma_{SFR}$  on metallicity over the entire range of gas surface densities in contrast to other models, and can also explain part of the scatter in the observations. We provide a tabulated form of the star formation laws that can be easily incorporated into numerical simulations or semi-analytical models of galaxy formation and evolution.

Accepted by Astrophysical Journal Letters

<http://arxiv.org/abs/1106.3282>

## Warm H<sub>2</sub>O and OH disk emission in V1331 Cyg

Greg W. Doppmann<sup>1</sup>, Joan R. Najita<sup>1</sup>, John S. Carr<sup>2</sup> and James R. Graham<sup>3,4</sup>

<sup>1</sup> National Optical Astronomy Observatory, USA

<sup>2</sup> Naval Research Laboratory, USA

<sup>3</sup> Astronomy Department, UC Berkeley, USA

<sup>4</sup> Dunlap Institute for Astronomy and Astrophysics, USA

E-mail contact: [gdoppmann@noao.edu](mailto:gdoppmann@noao.edu)

We present high resolution ( $R = 24,000$ )  $L$ -band spectra of the young intermediate mass star V1331 Cyg obtained with NIRSPEC on the Keck II telescope. The spectra show strong, rich emission from water and OH that likely arises from the warm surface region of the circumstellar disk. We explore the use of the new BT2 (Barber et al. 2006) water line list in fitting the spectra, and we find that it does a much better job than the well-known HITRAN (Rothman et al. 1998) water line list in the observed wavelength range and for the warm temperatures probed by our data. By comparing the observed spectra with synthetic disk emission models, we find that the water and OH emission lines have similar widths ( $\text{FWHM} \simeq 18 \text{ km s}^{-1}$ ). If the line widths are set by disk rotation, the OH and water emission lines probe a similar range of disk radii in this source. The water and OH emission are consistent with thermal emission for both components at a temperature  $\sim 1500 \text{ K}$ . The column densities of the emitting water and OH are large,  $\sim 10^{21} \text{ cm}^{-2}$  and  $\sim 10^{20} \text{ cm}^{-2}$ , respectively. Such a high column density of water is more than adequate to shield the disk midplane from external UV irradiation in the event of complete dust settling out of the disk atmosphere, enabling chemical synthesis to continue in the midplane despite a harsh external UV environment. The large OH-to-water ratio is similar to expectations for UV irradiated disks (e.g., Bethell and Bergin 2009), although the large OH column density is less easily accounted for.

Accepted by ApJ

<http://arxiv.org/abs/1106.5173>

# Multi-wavelength modeling of the spatially resolved debris disk of HD 107146

Steve Ertel<sup>1</sup>, Sebastian Wolf<sup>1</sup>, Stanimir Metchev<sup>2</sup>, Glenn Schneider<sup>3</sup>, John M. Carpenter<sup>4</sup>, Michael R. Meyer<sup>5</sup>, Lynne A. Hillenbrand<sup>4</sup> and Murray D. Silverstone<sup>6</sup>

<sup>1</sup> Institute of Theoretical Physics und Astrophysics, University of Kiel, Leibnizstraße 15, 24098 Kiel, Germany

<sup>2</sup> Department of Physics and Astronomy, State University of New York, Stony Brook, NY 11794-3800, USA

<sup>3</sup> Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721, USA

<sup>4</sup> Department of Astronomy, California Institute of Technology, 1200 East California Boulevard, Pasadena, CA 91125, USA

<sup>5</sup> Institute for Astronomy, ETH Zürich, Wolfgang-Pauli-Straße 27, 8093 Zürich, Switzerland

<sup>6</sup> Department of Physics & Astronomy, University of Alabama, 326 Gallalee, Tuscaloosa, AL 35487-0324, USA

E-mail contact: sertil *at* astrophysik.uni-kiel.de

*Aims:* We aim to constrain the location, composition, and dynamical state of planetesimal populations and dust around the young, sun-like (G2 V) star HD 107146.

*Methods:* We consider coronagraphic observations obtained with the Advanced Camera for Surveys (*HST*/ACS) onboard the *Hubble Space Telescope* (*HST*) in broad V ( $\lambda_c \approx 0.6 \mu\text{m}$ ) and broad I ( $\lambda_c \approx 0.8 \mu\text{m}$ ) filters, a resolved 1.3 mm map obtained with the *Combined Array for Research in Millimeter-Wave Astronomy* (*CARMA*), *Spitzer*/IRS low resolution spectra in the range of  $7.6 \mu\text{m}$  to  $37.0 \mu\text{m}$ , and the spectral energy distribution (SED) of the object at wavelengths ranging from  $3.5 \mu\text{m}$  to 3.1 mm. We complement these data with new coronagraphic high resolution observations of the debris disk using the Near Infrared Camera and Multi-Object Spectrometer (*HST*/NICMOS) aboard the *HST* in the F110W filter ( $\lambda_c \approx 1.1 \mu\text{m}$ ). The SED and images of the disk in scattered light as well as in thermal reemission are combined in our modeling using a parameterized model for the disk density distribution and optical properties of the dust.

*Results:* A detailed analytical model of the debris disk around HD 107146 is presented that allows us to reproduce the almost entire set of spatially resolved and unresolved multi-wavelength observations. Considering the variety of complementary observational data, we are able to break the degeneracies produced by modeling SED data alone. We find the disk to be an extended ring with a peak surface density at 131 AU. Furthermore, we find evidence for an additional, inner disk probably composed of small grains released at the inner edge of the outer disk and moving inwards due to Poynting-Robertson drag. A birth ring scenario (i.e., a more or less broad ring of planetesimals creating the dust disk through collisions) is found to be the most likely explanation of the ringlike shape of the disk.

Accepted by Astronomy and Astrophysics

<http://arxiv.org/abs/1107.1057>

## A rotating molecular disk toward IRAS 18162-2048, the exciting source of HH 80-81

Manuel Fernández-López<sup>1,6</sup>, Josep Miquel Girart<sup>2</sup>, Salvador Curiel<sup>1</sup>, Yolanda Gómez<sup>3</sup>, Paul T. P. Ho<sup>4,5</sup> and Nimesh Patel<sup>5</sup>

<sup>1</sup> Instituto de Astronomía, Universidad Nacional Autónoma de México (UNAM), Apartado Postal 70-264, 04510 Mexico, DF, Mexico

<sup>2</sup> Institut de Ciències de l'Espai, (CSIC-IEEC), Campus UAB, Facultat de Ciències, Torre C5-parell 2, 08193 Bellaterra, Catalunya, Spain

<sup>3</sup> Centro de Radioastronomía y Astrofísica, UNAM, Apartado Postal 3-72, Morelia, Michoacán 58089, México

<sup>4</sup> Academia Sinica Institute of Astronomy and Astrophysics, P.O. Box 23-141, Taipei 10617, Taiwan

<sup>5</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

<sup>6</sup> Current address: Department of Astronomy, University of Illinois at Urbana-Champaign, 1002 West Green Street, Urbana, IL 61801, USA

E-mail contact: manferna *at* gmail.com

We present several molecular line emission arcsec and subarcsec observations obtained with the Submillimeter Array (SMA) in the direction of the massive protostar IRAS 18162–2048, the exciting source of HH 80–81.

The data clearly indicates the presence of a compact (radius~425–850 AU) SO<sub>2</sub> structure, enveloping the more compact (radius<150 AU) 1.4 millimeter dust emission (reported in a previous paper). The emission spatially coincides with the position of the prominent thermal radio jet which terminates at the HH 80–81 and HH 80N Herbig–Haro objects.

Furthermore, the molecular emission is elongated in the direction perpendicular to the axis of the thermal radio jet, suggesting a disk-like structure. We derive a total dynamic mass (disk-like structure and protostar) of 11–15  $M_{\odot}$ . The  $\text{SO}_2$  spectral line data also allow us to constrain the structure temperature between 120–160 K and the volume density  $> 2 \times 10^9 \text{ cm}^{-3}$ . We also find that such a rotating flattened system could be unstable due to gravitational disturbances.

The data from  $\text{C}^{17}\text{O}$  line emission show a dense core within this star-forming region. Additionally, the  $\text{H}_2\text{CO}$  and the  $\text{SO}$  emissions appear clumpy and trace the disk-like structure, a possible interaction between a molecular core and the outflows, and in part, the cavity walls excavated by the thermal radio jet.

Accepted by AJ, 07-15-2011

## Unveiling Sources of Heating in the Vicinity of the Orion BN/KL Hot Core as Traced by Highly Excited Inversion Transitions of Ammonia

C. Goddi<sup>1,2</sup>, L. J. Greenhill<sup>2</sup>, E. M. L. Humphreys<sup>1,2</sup>, C. J. Chandler<sup>3</sup> and L. D. Matthews<sup>4</sup>

<sup>1</sup> European Southern Observatory, Karl-Schwarzschild-Strasse 2, D-85748 Garching bei Munchen, Germany

<sup>2</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

<sup>3</sup> National Radio Astronomy Observatory, P.O. Box O, Socorro, NM 87801, USA

<sup>4</sup> MIT Haystack Observatory, Westford, MA 01886, USA

E-mail contact: cgoddi at eso.org

Using the Expanded Very Large Array, we have mapped the vicinity of the Orion BN/KL Hot Core with sub-arcsecond angular resolution in seven metastable inversion transitions of ammonia ( $\text{NH}_3$ ): (J,K)=(6,6) to (12,12). This emission comes from levels up to 1500 K above the ground state, enabling identification of source(s) responsible for heating the region. We used this multi-transition dataset to produce images of the rotational/kinetic temperature ( $T_{\text{rot}}/T_{\text{kin}}$ ) and the column density  $N_{\text{col}}$  of  $\text{NH}_3$  for ortho and para species separately and on a position-by-position basis. We find  $T_{\text{rot}}$  and  $N_{\text{col}}$  in the range 160–490 K and  $(1-4) \times 10^{17} \text{ cm}^{-2}$ , respectively. Our spatially-resolved images show that the highest (column) density and hottest gas is found in a northeast-southwest elongated ridge to the southeast of Source I. We have also measured the ortho-para ratio of ammonia, estimated to vary in the range 0.9-1.6. Enhancement of ortho with respect to para and the offset of hot  $\text{NH}_3$  emission peaks from known (proto)stellar sources provide evidence that the  $\text{NH}_3$  molecules have been released from dust grains into the gas-phase through the passage of shocks and not by stellar radiation. We propose that the combined effect of Source I's proper motion and its low-velocity outflow impinging on a pre-existing dense medium is responsible for the excitation of  $\text{NH}_3$  and the Orion Hot Core. Finally, we found for the first time evidence of a slow ( $\sim 5 \text{ km s}^{-1}$ ) and compact ( $\sim 1000 \text{ AU}$ ) outflow towards IRC7.

Accepted by ApJ Letters (EVLA Special Issue)

<http://arxiv.org/abs/1106.4202>

## Mid-infrared interferometry of the massive young stellar object NGC 2264 IRS 1

R. Grellmann<sup>1</sup>, Th. Ratzka<sup>1</sup>, S. Kraus<sup>2</sup>, H. Linz<sup>3</sup>, Th. Preibisch<sup>1</sup> and G. Weigelt<sup>4</sup>

<sup>1</sup> Universitaets-Sternwarte, Ludwig-Maximilians-Universitaet Muenchen, Scheinerstr. 1, 81679 Muenchen, Germany

<sup>2</sup> University of Michigan, Department of Astronomy, Ann Arbor, MI 48109-1090, USA

<sup>3</sup> Max-Planck-Institut fuer Astronomie, Koenigstuhl 17, 69117 Heidelberg, Germany

<sup>4</sup> Max-Planck-Institut fuer Radioastronomie, Auf dem Huegel 69, 53121 Bonn, Germany

E-mail contact: grellman at usm.uni-muenchen.de

The optically invisible infrared-source NGC 2264 IRS 1 lying north of the Cone Nebula is thought to be a massive young stellar object ( $\sim 10 M_{\text{sun}}$ ). Although strong infrared excess clearly shows that the central object is surrounded by large amounts of circumstellar material, no information about the spatial distribution of this circumstellar material has been available until now. We used the ESO Very Large Telescope Interferometer to perform long-baseline interferometric observations of NGC 2264 IRS 1 in the mid-infrared regime. Our observations resolve the circumstellar material around NGC 2264 IRS 1, provide the first direct measurement of the angular size of the mid-infrared emission, and yield direct constraints on the spatial distribution of the dust. We analyze the spectrally dispersed interferometric data taken with MIDI at two different position angles and baseline lengths. We use different approaches (a geometrical model, a

temperature-gradient model, and radiative transfer models) to jointly model the observed interferometric visibilities and the spectral energy distribution. The derived visibility values between 0.02 and 0.3 show that the mid-infrared emission is clearly resolved. The characteristic size of the MIR-emission region is 30-60 AU; this value is typical for other YSOs with similar or somewhat lower luminosities. A comparison of the sizes for the two position angles shows a significant elongation of the dust distribution. Simple spherical envelope models are therefore inconsistent with the data. The radiative transfer modeling of our data suggests that we observe a geometrically thin and optically thick circumstellar disk with a mass of about 0.1 Msun. Our modeling indicates that NGC 2264 IRS 1 is surrounded by a flat circumstellar disk that has properties similar to disks typically found around lower-mass young stellar objects. This result supports the assumption that massive young stellar objects form via accretion from circumstellar disks.

Accepted by A&A

## **A Correlation Between Surface Densities of Young Stellar Objects and Gas in Eight Nearby Molecular Clouds**

**R. A. Gutermuth<sup>1,2</sup>, J. L. Pipher<sup>3</sup>, S. T. Megeath<sup>4</sup>, P. C. Myers<sup>5</sup>, L. E. Allen<sup>6</sup>, & T. S. Allen<sup>4</sup>**

<sup>1</sup> Five College Astronomy Dept., Smith College, Northampton, MA 01063

<sup>2</sup> Dept. of Astronomy, University of Massachusetts, Amherst, MA

<sup>3</sup> Dept. of Physics & Astronomy, University of Rochester, Rochester, NY

<sup>4</sup> Dept. of Physics & Astronomy, University of Toledo, Toledo, OH

<sup>5</sup> Harvard-Smithsonian Center for Astrophysics, Cambridge, MA

<sup>6</sup> National Optical Astronomy Observatories, Tucson, AZ

E-mail contact: rgutermu *at* astro.umass.edu

We report the discovery and characterization of a power law correlation between the local surface densities of Spitzer-identified, dusty young stellar objects and the column density of gas (as traced by near-IR extinction) in eight molecular clouds within 1 kpc and with 100 or more known YSOs. This correlation, which appears in data smoothed over size scales of  $\sim 1$  pc, varies in quality from cloud to cloud; those clouds with tight correlations, MonR2 and Ophiuchus, are fit with power laws of slope 2.67 and 1.87, respectively. The spread in the correlation is attributed primarily to local gas disruption by stars that formed there or to the presence of very young sub-regions at the onset of star formation. We explore the ratio of the number of Class II to Class I sources, a proxy for the star formation age of a region, as a function of gas column density; this analysis reveals a declining Class II to Class I ratio with increasing column density. We show that the observed star-gas correlation is consistent with a star formation law where the star formation rate per area varies with the gas column density squared. We also propose a simple picture of thermal fragmentation of dense gas in an isothermal, self-gravitating layer as an explanation for the power law. Finally, we briefly compare the star gas correlation and its implied star formation law with other recent proposed of star formation laws at similar and larger size scales from nearby star forming regions.

Accepted by ApJ

## **Resolved Submillimeter Observations of the HR 8799 and HD 107146 Debris Disks**

**A. Meredith Hughes<sup>1</sup>, David J. Wilner<sup>2</sup>, Sean M. Andrews<sup>2</sup>, Jonathan P. Williams<sup>3</sup>, Kate Y. L. Su<sup>4</sup>, Ruth A. Murray-Clay<sup>2</sup> and Chunhua Qi<sup>2</sup>**

<sup>1</sup> Miller Fellow, UC Berkeley Dept. of Astronomy, 601 Campbell Hall, Berkeley, CA 94720, USA

<sup>2</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138, USA

<sup>3</sup> Institute for Astronomy, University of Hawaii, 2680 Woodlawn Dr., Honolulu, HI 96822, USA

<sup>4</sup> Steward Observatory, University of Arizona, 933 N. Cherry Ave, Tucson, AZ 85721, USA

E-mail contact: mhughes *at* astro.berkeley.edu

We present 880  $\mu\text{m}$  Submillimeter Array observations of the debris disks around the young solar analogue HD 107146 and the multiple-planet host star HR 8799, at an angular resolution of 3" and 6", respectively. We spatially resolve the inner edge of the disk around HR 8799 for the first time. While the data are not sensitive enough (with rms noise of 1 mJy) to constrain the system geometry, we demonstrate that a model by Su et al. (2009) based on the spectral energy distribution (SED) with an inner radius of 150 AU predicts well the spatially resolved data. Furthermore, by

modeling simultaneously the SED and visibilities, we demonstrate that the dust is distributed in a broad (of order 100 AU) annulus rather than a narrow ring. We also model the observed SED and visibilities for the HD 107146 debris disk and generate a model of the dust emission that extends in a broad band between 50 and 170 AU from the star. We perform an *a posteriori* comparison with existing 1.3 mm CARMA observations and demonstrate that a smooth, axisymmetric model reproduces well all of the available millimeter-wavelength data.

Accepted by The Astrophysical Journal

[http://astro.berkeley.edu/~mhughes/download/hughes\\_debris.pdf](http://astro.berkeley.edu/~mhughes/download/hughes_debris.pdf)

## Variability of the SiO thermal line emission toward the young L1448-mm outflow.

Izaskun Jimenez-Serra<sup>1</sup>, Jesus Martin-Pintado<sup>2</sup>, Jan-Martin Winters<sup>3</sup>, Arturo Rodriguez-Franco<sup>2</sup> and Paola Caselli<sup>4</sup>

<sup>1</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138, USA

<sup>2</sup> Centro de Astrobiología (CSIC/INTA), Ctra. de Torrejón a Ajalvir km 4, E-28850 Torrejón de Ardoz, Madrid, Spain

<sup>3</sup> Institut de Radio Astronomie Millimétrique, 300 Rue de la Piscine, F-38406 St. Martin d'Hères, France

<sup>4</sup> School of Physics & Astronomy, E.C. Stoner Building, The University of Leeds, Leeds, LS2 9JT, UK

E-mail contact: [ijimenez-serra at cfa.harvard.edu](mailto:ijimenez-serra@cfa.harvard.edu)

The detection of narrow SiO thermal emission toward young outflows has been proposed to be a signature of the magnetic precursor of C-shocks. Recent modeling of the SiO emission across C-shocks predicts variations in the SiO line intensity and line shape at the precursor and intermediate-velocity regimes in only few years. We present high-angular resolution ( $3.8'' \times 3.3''$ ) images of the thermal SiO  $J=2 \rightarrow 1$  emission toward the L1448-mm outflow in two epochs (November 2004-February 2005, March-April 2009). Several SiO condensations have appeared at intermediate velocities ( $20\text{-}50 \text{ km s}^{-1}$ ) toward the red-shifted lobe of the outflow since 2005. Toward one of the condensations (clump D), systematic differences of the dirty beams between 2005 and 2009 could be responsible for the SiO variability. At higher velocities ( $50\text{-}80 \text{ km s}^{-1}$ ), SiO could also have experienced changes in its intensity. We propose that the SiO variability toward L1448-mm is due to a real SiO enhancement by young C-shocks at the internal working surface between the jet and the ambient gas. For the precursor regime ( $5.2\text{-}9.2 \text{ km s}^{-1}$ ), several narrow and faint SiO components are detected. Narrow SiO tends to be compact, transient and shows elongated (bow-shock) morphologies perpendicular to the jet. We speculate that these features are associated with the precursor of C-shocks appearing at the interface of the new SiO components seen at intermediate velocities.

Accepted by Astrophysical Journal

<http://arxiv.org/abs/1107.2067>

## Confirmation of the VeLLO L1148-IRS: Star Formation at very low (Column) Density

Jens Kauffmann<sup>1</sup>, Frank Bertoldi<sup>2</sup>, Tyler L. Bourke<sup>3</sup>, Philip C. Myers<sup>3</sup>, Chang Won Lee<sup>4</sup> and Tracy L. Huard<sup>5</sup>

<sup>1</sup> NPP Fellow, Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 91109, USA

<sup>2</sup> Argelander Institut fuer Astronomie, Universitaet Bonn, Auf dem Huegel 71, 53121 Bonn, Germany

<sup>3</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, MA 02138, USA

<sup>4</sup> Korea Astronomy and Space Science Institute, 61-1 Hwaam-dong, Yusung-gu, Daejeon 305-348, Korea

<sup>5</sup> Department of Astronomy, University of Maryland, College Park, MD 20742, USA

E-mail contact: [jens.kauffmann at jpl.nasa.gov](mailto:jens.kauffmann@jpl.nasa.gov)

We report the detection of a compact ( $\sim 5''$ ; about 1800 AU projected size) CO outflow from L1148-IRS. This confirms that this Spitzer source is physically associated with the nearby ( $\approx 325 \text{ pc}$ ) L1148 dense core. Radiative transfer modeling suggests an internal luminosity of 0.08 to  $0.13 L_{\odot}$ . This validates L1148-IRS as a Very Low Luminosity Object (VeLLO;  $L \leq 0.1 L_{\odot}$ ). The L1148 dense core has unusually low densities and column densities for a star-forming core. It is difficult to understand how L1148-IRS might have formed under these conditions. Independent of the exact final mass of this VeLLO (which is likely  $< 0.24 M_{\odot}$ ), L1148-IRS and similar VeLLOs might hold some clues about the isolated formation of brown dwarfs.

Accepted by MNRAS

<http://arxiv.org/abs/1106.2545>

## Dynamical Friction in a Gas: The Subsonic Case

Aaron T. Lee<sup>1</sup> and Steven W. Stahler<sup>1</sup>

<sup>1</sup> Astronomy Dept, U. of California, Berkeley, CA 94720 USA

E-mail contact: a.t.lee *at* berkeley.edu

We study the force of dynamical friction acting on a gravitating point mass that travels through an extended, isothermal gas. This force is well established in the hypersonic limit, but remains less understood in the subsonic regime. Using perturbation theory, we analyze the changes in gas velocity and density far from the mass. We show analytically that the steady-state friction force is  $\dot{M}V$ , where  $\dot{M}$  is the mass accretion rate onto an object moving at speed  $V$ . It follows that the speed of an object experiencing no other forces declines as the inverse square of its mass. Using a modified version of the classic Bondi-Hoyle interpolation formula for  $\dot{M}$  as a function of  $V$ , we derive an analytic expression for the friction force. This expression also holds when mass accretion is thwarted, e.g. by a wind, as long as the wind-cloud interaction is sufficiently confined spatially. Our result should find application in a number of astrophysical settings, such as the motion of galaxies through intracluster gas.

Accepted by MNRAS

<http://arxiv.org/abs/1106.4820>

## Massive star formation around I05345+3157 – I. The dense gas

Katherine I. Lee<sup>1</sup>, Leslie W. Looney<sup>1</sup>, Randolph Klein<sup>2,3</sup> and Shiya Wang<sup>4</sup>

<sup>1</sup> Department of Astronomy, University of Illinois at Urbana-Champaign, 1002 W Green St, Urbana IL 61801, USA

<sup>2</sup> SOFIA-USRA, NASA Ames Research Center, Mail Stop N211-3, Moffett Field, CA 94035, USA

<sup>3</sup> Department of Physics, University of California at Berkeley, 366 Le Conte Hall, Berkeley, CA 94720, USA

<sup>4</sup> Department of Astronomy, University of Michigan at Ann Arbor, 500 Church St., Ann Arbor, MI 48109, USA

E-mail contact: ijlee9 *at* illinois.edu

We present observations of the intermediate to massive star-forming region I05345+3157 using the molecular line tracer CS(2–1) with CARMA to reveal the properties of the dense gas cores. Seven gas cores are identified in the integrated intensity map of CS(2–1). Among these, core 1 and core 3 have counterparts in the  $\lambda = 2.7$  mm continuum data. We suggest that core 1 and core 3 are star-forming cores that may already or will very soon harbor young massive protostars. The total masses of core 1 estimated from the LTE method and dust emission by assuming a gas-to-dust ratio are  $5 \pm 1 M_{\odot}$  and  $18 \pm 6 M_{\odot}$ , and that of core 3 are  $15 \pm 7 M_{\odot}$  and  $11 \pm 3 M_{\odot}$ . The spectrum of core 3 shows blue-skewed self-absorption, which suggests gas infall – a collapsing core. The observed broad linewidths of the seven gas cores indicate non-thermal motions. These non-thermal motions can be interactions with nearby outflows or due to the initial turbulence; the former is observed, while the role of initial turbulence is less certain. Finally, the virial masses of the gas cores are larger than the LTE masses, which for a bound core implies a requirement on the external pressure of  $\sim 10^8 \text{ K cm}^{-3}$ . The cores have the potential to further form massive stars.

Accepted by MNRAS

<http://arxiv.org/abs/1107.1224>

## Clustered Star Formation in Magnetic Clouds: Properties of Dense Cores Formed in Outflow-Driven Turbulence

Fumitaka Nakamura<sup>1</sup> and Zhi-Yun Li<sup>2</sup>

<sup>1</sup> National Astronomical Observatory of Japan

<sup>2</sup> University of Virginia

E-mail contact: fumitaka.nakamura *at* nao.ac.jp

We investigate the physical properties of dense cores formed in turbulent, magnetized, parsec-scale clumps of molecular clouds, using three-dimensional numerical simulations that include protostellar outflow feedback. The dense cores are

identified in the simulated density data cube through a clumpfind algorithm. We find that the core velocity dispersion does not show any clear dependence on the core size, in contrast to Larson's linewidth-size relation, but consistent with recent observations. In the absence of a magnetic field, the majority of the cores have supersonic velocity dispersions. A moderately-strong magnetic field reduces the dispersion to a subsonic or at most transonic value typically. Most of the cores are out of virial equilibrium, with the external pressure dominating the self-gravity. The implication is that the core evolution is largely controlled by the outflow-driven turbulence. Even an initially-weak magnetic field can retard star formation significantly, because the field is amplified by the outflow-driven turbulence to an equipartition strength, with the distorted field component dominating the uniform one. In contrast, for a moderately-strong field, the uniform component remains dominant. Such a difference in the magnetic structure is evident in our simulated polarization maps of dust thermal emission; it provides a handle on the field strength. Recent polarization measurements show that the field lines in cluster-forming clumps are spatially well-ordered. It is indicative of a moderately-strong, dynamically important, field which, in combination with outflow feedback, can keep the rate of star formation in embedded clusters at the observationally-inferred, relatively-slow rate of several percent per free-fall time.

Accepted by Astrophysical Journal

## **Search of HH Objects and Emission-Line Stars in Star Forming Regions. VII. Herbig-Haro Objects in the Vicinity of GM 2-41 Nebula.**

**E. H. Nikogosyan<sup>1</sup>, T. Yu. Magakian<sup>1</sup> and T. A. Movsessian<sup>1</sup>**

<sup>1</sup> Byurakan Astrophysical Observatory, Aragatsotn prov., Armenia

E-mail contact: elena *at* bao.sci.am

Five new HH objects (HH 1036 -1040) were revealed in the area 14 x 14 arcmin in the vicinity of GM 2-41 nebula in the central part of the HII region DR 15, located in the southern part of the Cyg OB2 association. Four of them have complex structure, resembling the HH-flows. In HH 1036 on the archival images of Spitzer database the infrared emission of molecular hydrogen was detected. Also two new infrared nebulae, illuminated by very red young stellar objects, were found.

Accepted by Astrophysics

## **Search of HH Objects and Emission-Line Stars in Star Forming Regions. VIII. H $\alpha$ emission stars around the GM 2-41 Nebula.**

**E. H. Nikogosyan<sup>1</sup>, T. Yu. Magakian<sup>1</sup> and T. A. Movsessian<sup>1</sup>**

<sup>1</sup> Byurakan Astrophysical Observatory, Aragatsotn prov., Armenia

E-mail contact: elena *at* bao.sci.am

43 stars with H $\alpha$  emission in the area 14 x 14 arcmin around the GM 2-41 nebula, located in the HII region DR 15 in the southern part of the Cyg OB2 association, were revealed by the method of slitless spectroscopy. 30 objects are new ones. On the basis of VI and JHK photometric data the majority of these objects are classified as low-mass PMS objects with F7 - M3 spectral type and about 1 Myr age. The distribution of the emission stars is not uniform. They form two main groups which by coordinates coincide with the UCHII G0.79+0.3 and G79.2+0.4 regions.

Accepted by Astrophysics

## **A highly efficient measure of mass segregation in star clusters**

**Christoph Olczak<sup>1,2,3,4</sup>, Rainer Spurzem<sup>3,1,4</sup> and Thomas Henning<sup>2</sup>**

<sup>1</sup> Astronomisches Rechen-Institut (ARI), Zentrum für Astronomie Universität Heidelberg, Mönchhofstrasse 12-14, 69120 Heidelberg

<sup>2</sup> Max-Planck-Institut für Astronomie (MPIA), Königstuhl 17, 69117 Heidelberg, Germany

<sup>3</sup> National Astronomical Observatories of China, Chinese Academy of Sciences (NAOC/CAS), 20A Datun Lu, Chaoyang District, Beijing 100012, China

<sup>4</sup> The Kavli Institute for Astronomy and Astrophysics at Peking University (KIAA), Yi He Yuan Lu 5, Hai Dian Qu, Beijing 100871, China



E-mail contact: *olczak at ari.uni-heidelberg.de*

Investigations of mass segregation are of vital interest for the understanding of the formation and dynamical evolution of stellar systems on a wide range of spatial scales. Our method is based on the minimum spanning tree (MST) that serves as a geometry-independent measure of concentration. Compared to previous such approaches we obtain a significant refinement by using the geometrical mean as an intermediate-pass. It allows the detection of mass segregation with much higher confidence and for much lower degrees of mass segregation than other approaches. The method shows in particular very clear signatures even when applied to small subsets of the entire population. We confirm with high significance strong mass segregation of the five most massive stars in the Orion Nebula Cluster (ONC). Our method is the most sensitive general measure of mass segregation so far and provides robust results for both data from simulations and observations. As such it is ideally suited for tracking mass segregation in young star clusters and to investigate the long standing paradigm of primordial mass segregation by comparison of simulations and observations.

Accepted by A&A

<http://arxiv.org/abs/1107.1842>

## Numerical convergence in self-gravitating disc simulations: initial conditions and edge effects

Sijme-Jan Paardekooper<sup>1</sup>, Clément Baruteau<sup>1</sup> and Farzana Meru<sup>2</sup>

<sup>1</sup> DAMTP, University of Cambridge, Wilberforce Road, Cambridge CB3 0WA, UK

<sup>2</sup> Institut für Astronomie and Astrophysik, Universität Tübingen, Auf der Morgenstelle 10, Tübingen D-72076, Germany

E-mail contact: *S.Paardekooper at damtp.cam.ac.uk*

We study the numerical convergence of hydrodynamical simulations of self-gravitating accretion discs, in which a simple cooling law is balanced by shock heating. It is well-known that there exists a critical cooling time scale for which shock heating can no longer compensate for the energy losses, at which point the disc fragments. The numerical convergence of previous results of this critical cooling time scale was questioned recently using Smoothed Particle Hydrodynamics (SPH). We employ a two-dimensional grid-based code to study this problem, and find that for smooth initial conditions, fragmentation is possible for slower cooling as the resolution is increased, in agreement with recent SPH results. We show that this non-convergence is at least partly due to the creation of a special location in the disc, the boundary between the turbulent and the laminar region, when cooling towards a gravito-turbulent state. Converged results appear to be obtained in setups where no such sharp edges appear, and we then find a critical cooling time scale of  $\sim 4 \Omega^{-1}$ , where  $\Omega$  is the local angular velocity.

Accepted by MNRAS

<http://arxiv.org/abs/1106.4425>

## Probing the initial conditions of high-mass star formation — II. Fragmentation, stability, and chemistry towards high-mass star-forming regions G29.96–0.02 and G35.20–1.74

T. Pillai<sup>1,2</sup>, J. Kauffmann<sup>2,3,4</sup>, F. Wyrowski<sup>5</sup>, J. Hatchell<sup>6</sup>, A.G. Gibb<sup>7</sup>, M.A. Thompson<sup>8</sup>

<sup>1</sup> California Institute Of Technology, 1200 E California Blvd, Pasadena, CA 91125, USA

<sup>2</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

<sup>3</sup> Initiative in Innovative Computing (IIC), Harvard University, 60 Oxford Street, Cambridge, MA 02138, USA

<sup>4</sup> Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 91109, USA

<sup>5</sup> Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany

<sup>6</sup> University of Exeter, UK

<sup>7</sup> Department of Physics & Astronomy, University of British Columbia, Vancouver, BC, V6T 1Z1, Canada

<sup>8</sup> Centre for Astrophysics Research, Science & Technology Research Institute, University of Hertfordshire, College Lane, Hatfield, AL10 9AB, UK

E-mail contact: *tpillai at astro.caltech.edu*

Most work on high-mass star formation has focused on observations of young massive stars in protoclusters. Very little is known about the preceding stage. Here, we present a new high-resolution study of pre-protocluster regions in tracers exclusively probing the coldest and dense gas ( $\text{NH}_2\text{D}$ ). The two target regions G29.96–0.02 and G35.20–1.74 (W48) are drawn from the SCAMPS project, which searches for pre-protoclusters near known ultracompact HII regions. We used our data to constrain the chemical, thermal, kinematic, and physical conditions (i.e., densities) in G29.96e and G35.20w.  $\text{NH}_3$ ,  $\text{NH}_2\text{D}$ ,  $\text{HCO}^+$ , and continuum emission were mapped using the VLA, PdBI, and BIMA. In particular,  $\text{NH}_2\text{D}$  is a unique tracer of cold, precluster gas at high densities, while  $\text{NH}_3$  traces both the cold and warm gas of modest-to-high densities. In G29.96e, Spitzer images reveal two massive filaments, one of them in extinction (infrared dark cloud). Dust and line observations reveal fragmentation into multiple massive cores strung along filamentary structures. Most of these are cold ( $< 20$  K), dense ( $> 10^5 \text{ cm}^{-3}$ ) and highly deuterated ( $[\text{NH}_2\text{D}/\text{NH}_3] > 6\%$ ). In particular, we observe very low line widths in  $\text{NH}_2\text{D}$  ( $\text{FWHM} \leq 1 \text{ km s}^{-1}$ ). These are very narrow lines that are unexpected towards a region forming massive stars. Only one core in the center of each filament appears to be forming massive stars (identified by the presence of masers and massive outflows); however, it appears that only a few such stars are currently forming (i.e., just a single Spitzer source per region). These multi-wavelength, high-resolution observations of high-mass pre-protocluster regions show that the target regions are characterized by (i) turbulent Jeans fragmentation of massive clumps into cores (from a Jeans analysis); (ii) cores and clumps that are “over-bound/subvirial”, i.e. turbulence is too weak to support them against collapse, meaning that (iii) some models of monolithic cloud collapse are quantitatively inconsistent with data; (iv) accretion from the core onto a massive star, which can (for observed core sizes and velocities) be sustained by accretion of envelope material onto the core, suggesting that (similar to competitive accretion scenarios) the mass reservoir for star formation is not necessarily limited to the natal core; (v) high deuteration ratios ( $[\text{NH}_2\text{D}/\text{NH}_3] > 6\%$ ), which make the above discoveries possible; (vi) and the destruction of  $\text{NH}_2\text{D}$  toward embedded stars.

Accepted by Astronomy & Astrophysics, Volume 530, article 118

<http://adsabs.harvard.edu/abs/2011A%26A...530A.118P>

## EVLA Observations of the Barnard 5 Star-Forming Core: Embedded Filaments Revealed

Jaime E Pineda<sup>1,2</sup>, Alyssa A. Goodman<sup>3</sup>, Hector Arce<sup>4</sup>, Paola Caselli<sup>5</sup>, Steven Longmore<sup>1</sup> and Stuartt Corder<sup>6,7</sup>

<sup>1</sup> European Southern Observatory, Karl Schwarzschild Str. 2, 85748 Garching bei Munchen, Germany

<sup>2</sup> UK ALMA Regional Centre Node, Jodrell Bank Centre for Astrophysics, School of Physics and Astronomy, University of Manchester, Manchester, M13 9PL, UK

<sup>3</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138, USA

<sup>4</sup> Department of Astronomy, Yale University, P.O. Box 208101, New Haven, CT 06520-8101, USA

<sup>5</sup> School of Physics and Astronomy, University of Leeds, Leeds LS2 9JT, UK

<sup>6</sup> North American ALMA Science Center, 520 Edgemont Road, Charlottesville, VA 22903, USA

<sup>7</sup> National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, VA 22903, USA

E-mail contact: Jaime.Pineda *at* manchester.ac.uk

We present a  $\sim 6.5' \times 8'$  Expanded Very Large Array (EVLA) mosaic observations of the  $\text{NH}_3(1,1)$  emission in the Barnard 5 region in Perseus, with an angular resolution of 6 arcsec. This map covers the coherent region, where the dense gas presents subsonic non-thermal motions (as seen from single dish observations with the Green Bank Telescope, GBT). The combined EVLA and GBT observations reveal, for the first time, a striking filamentary structure (20 arcsec wide or 5,000 AU at the distance of Perseus) in this low-mass star forming region. The integrated intensity profile of this structure is consistent with models of an isothermal filament in hydrostatic equilibrium. The observed separation between the B5–IRS1 young stellar object (YSO), in the central region of the core, and the northern starless condensation matches the Jeans length of the dense gas. This suggests that the dense gas in the coherent region is fragmenting. The region observed displays a narrow velocity dispersion, where most of the gas shows evidence for subsonic turbulence, and where little spatial variations are present. It is only close to the YSO where an increase in the velocity dispersion is found, but still displaying subsonic non-thermal motions.

Accepted by The Astrophysical Journal Letters (special issue on EVLA Early Science)

<http://arxiv.org/abs/1106.5474>

# Searching for Gas Giant Planets on Solar System Scales: VLT NACO/APP Observations of the Debris Disk Host Stars HD172555 and HD115892

Sascha P. Quanz<sup>1</sup>, Matthew A. Kenworthy<sup>2</sup>, Michael R. Meyer<sup>1</sup>, Julien H. V. Girard<sup>3</sup> and Markus Kasper<sup>4</sup>

<sup>1</sup> Institute for Astronomy, ETH Zurich, Wolfgang-Pauli-Strasse 27, 8093 Zurich, Switzerland

<sup>2</sup> Sterrewacht Leiden, P.O. Box 9513, Niels Bohrweg 2, 2300 RA Leiden, The Netherlands

<sup>3</sup> European Southern Observatory, Alonso de Cordova 3107, Vitacura, Cassilla 19001, Santiago, Chile

<sup>4</sup> European Southern Observatory, Karl Schwarzschild Strasse, 2, 85748 Garching bei München, Germany

E-mail contact: quanz *at* astro.phys.ethz.ch

Using the APP coronagraph of VLT/NACO we searched for planetary mass companions around HD115892 and HD172555 in the thermal infrared at 4  $\mu\text{m}$ . Both objects harbor unusually luminous debris disks for their age and it has been suggested that small dust grains were produced recently in transient events (e.g., a collision) in these systems. Such a collision of planetesimals or protoplanets could have been dynamically triggered by yet unseen companions. We did not detect any companions in our images but derived the following detection limits: For both objects we would have detected companions with apparent magnitudes between  $\sim 13.2$ – $14.1$  mag at angular separations between  $0.4$ – $1.0''$  at the  $5\text{-}\sigma$  level. For HD115892 we were sensitive to companions with  $12.1$  mag even at  $0.3''$ . Using theoretical models these magnitudes are converted into mass limits. For HD115892 we would have detected objects with  $10$ – $15$   $M_{\text{Jup}}$  at angular separations between  $0.4$ – $1.0''$  ( $7$ – $18$  AU). At  $0.3''$  ( $\sim 5.5$  AU) the detection limit was  $\gtrsim 25$   $M_{\text{Jup}}$ . For HD172555 we reached detection limits between  $2$ – $3$   $M_{\text{Jup}}$  at separations between  $0.5$ – $1.0''$  ( $15$ – $29$  AU). At  $0.4''$  ( $\sim 11$  AU) the detection limit was  $\gtrsim 4$   $M_{\text{Jup}}$ . Despite the non-detections our data demonstrate the unprecedented contrast performance of NACO/APP in the thermal infrared at very small inner working angles and we show that our observations are mostly background limited at separations  $\gtrsim 0.5''$ .

Accepted by ApJL

<http://xxx.lanl.gov/abs/1106.4528>

## Variable jets with non-top hat ejection cross sections: a model for the knots of the HH 34 jet

A.C. Raga<sup>1</sup>, A. Noriega-Crespo<sup>2</sup>, P. Kajdic<sup>3</sup>, F. De Colle<sup>4</sup> and A. Esquivel<sup>1</sup>

<sup>1</sup> Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México

<sup>2</sup> Spitzer Science Center, California Institute of Technology, USA

<sup>3</sup> Instituto de Geofísica, Universidad Nacional Autónoma de México

<sup>4</sup> Astronomy and Astrophysics Department, University of California, USA

E-mail contact: raga *at* nucleares.unam.mx

We compute axisymmetric, single-sinusoidal mode variable ejection models with a non-top hat ejection velocity cross section. We find that for decreasing edge-to-centre velocity ratios one obtains internal working surfaces with progressively more extended bow shock wings. These wings produce [S II] emission which partially fills in the inter-knot regions in predicted intensity maps. We then compute 3-mode models (with parameters appropriate for the HH 34 jet), and compare predicted intensity maps with archival HST images of HH 34. We find that a model with a moderate edge-to-centre velocity ratio produces knot structures with morphologies and time-variabilities with clear similarities to the observations of HH 34.

Accepted by RMxAA

## An interpretive ballistic model for quasi-symmetric bipolar jet systems

A.C. Raga<sup>1</sup>, A. Noriega-Crespo<sup>2</sup>, J.C. Rodríguez-Ramírez<sup>1</sup>, V. Lora<sup>3</sup>, K.R. Stapelfeldt<sup>4</sup> and S.J. Carey<sup>2</sup>

<sup>1</sup> Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México

<sup>2</sup> Spitzer Science Center, California Institute of Technology, USA

<sup>3</sup> Astronomisches Rechen-Institut Zentrum für Astronomie der Universität Heidelberg, Germany

<sup>4</sup> Jet propulsion Laboratory, California Institute of Technology, USA

E-mail contact: raga *at* nucleares.unam.mx

We present an analytic, ballistic model for quasi-symmetric jet/counterjet systems, considering both the non-relativistic and the relativistic cases. The model considers the presence of ejection time and velocity asymmetries, which produce offsets between the positions of the knots in jet/counterjet pairs. A fit of the non-relativistic model predictions to observations of two quasi-symmetric HH outflows (HH 34 and HH 111) allows us to obtain the magnitudes of the ejection time and velocity asymmetries of these systems.

Accepted by RMxAA

## **New Young Star Candidates in the Taurus-Auriga Region as Selected from WISE**

**L. M. Rebull<sup>1</sup>, X. P. Koenig<sup>2</sup>, D. L. Padgett<sup>1</sup>, S. Terebey<sup>3</sup>, P. M. McGehee<sup>4</sup>, L. A. Hillenbrand<sup>5</sup>, G. Knapp<sup>6</sup>, D. Leisawitz<sup>2</sup>, W. Liu<sup>4</sup>, A. Noriega-Crespo<sup>1</sup>, M. Ressler<sup>7</sup>, K. R. Stapelfeldt<sup>7</sup>, S. Fajardo-Acosta<sup>4</sup> and A. Mainzer<sup>7</sup>**

<sup>1</sup> Spitzer Science Center, Caltech, Pasadena, CA, USA

<sup>2</sup> Goddard Space Flight Center, Greenbelt, MD, USA

<sup>3</sup> Cal State University, Los Angeles, USA

<sup>4</sup> Infrared Processing and Analysis Center, USA

<sup>5</sup> California Institute of Technology, USA

<sup>6</sup> Princeton University, USA

<sup>7</sup> Jet Propulsion Laboratory, USA

E-mail contact: rebull *at* ipac.caltech.edu

The Taurus Molecular Cloud subtends a large solid angle on the sky, in excess of 250 square degrees. The search for legitimate Taurus members to date has been limited by sky coverage as well as the challenge of distinguishing members from field interlopers. The Wide-field Infrared Survey Explorer (WISE) has recently observed the entire sky, and we take advantage of the opportunity to search for young stellar object (YSO) candidate Taurus members from a  $\sim 260$  square degree region designed to encompass previously-identified Taurus members. We use near- and mid-infrared colors to select objects with apparent infrared excesses and incorporate other catalogs of ancillary data to present: a list of rediscovered Taurus YSOs with infrared excesses (taken to be due to circumstellar disks), a list of rejected YSO candidates (largely galaxies), and a list of 94 surviving candidate new YSO-like Taurus members. There is likely to be contamination lingering in this candidate list, and follow-up spectra are warranted.

Accepted by ApJS

<http://web.ipac.caltech.edu/staff/rebull/research.html>

## **A natural formation scenario for misaligned and short-period eccentric extrasolar planets**

**Ingo Thies<sup>1</sup>, Pavel Kroupa<sup>1</sup>, Simon P. Goodwin<sup>2</sup>, Dimitris Stamatellos<sup>3</sup> and Anthony P. Whitworth<sup>3</sup>**

<sup>1</sup> Argelander-Institute for Astronomy, University of Bonn, Germany

<sup>2</sup> Department of Physics and Astronomy, University of Sheffield, Sheffield S3 7RH, UK

<sup>3</sup> School of Physics & Astronomy, Cardiff University, Cardiff CF24 3AA, UK

E-mail contact: ithies *at* astro.uni-bonn.de

Recent discoveries of strongly misaligned transiting exoplanets pose a challenge to the established planet formation theory which assumes planetary systems to form and evolve in isolation. However, the fact that the majority of stars actually do form in star clusters raises the question how isolated forming planetary systems really are. Besides radiative and tidal forces the presence of dense gas aggregates in star-forming regions are potential sources for perturbations to protoplanetary discs or systems. Here we show that subsequent capture of gas from large extended accretion envelopes onto a passing star with a typical circumstellar disc can tilt the disc plane to retrograde orientation, naturally explaining the formation of strongly inclined planetary systems. Furthermore, the inner disc regions may become denser, and thus more prone to speedy coagulation and planet formation. Pre-existing planetary systems are compressed by gas inflows leading to a natural occurrence of close-in misaligned hot Jupiters and short-period eccentric planets. The likelihood of such events mainly depends on the gas content of the cluster and is thus expected to be highest in the

youngest star clusters.

Accepted by MNRAS

<http://arxiv.org/abs/1107.2113>

## Chemical abundances in the protoplanetary disk LV 2 (Orion) – II: High dispersion VLT observations and microjet properties

Yiannis Tsamis<sup>1</sup> and Jeremy Walsh<sup>1</sup>

<sup>1</sup> European Southern Observatory, Garching, Germany

E-mail contact: [ytsamis@eso.org](mailto:ytsamis@eso.org)

Integral field spectroscopy of the LV 2 proplyd is presented taken with the VLT/FLAMES Argus array at an angular resolution of  $0.31 \times 0.31$  arcsec<sup>2</sup> and velocity resolutions down to  $2 \text{ km s}^{-1}$  per pixel. Following subtraction of the local M42 emission, the spectrum of LV 2 is isolated from the surrounding nebula. We measured the heliocentric velocities and widths of a number of lines detected in the intrinsic spectrum of the proplyd, as well as in the adjacent Orion nebula falling within a  $6.6 \times 4.2$  arcsec<sup>2</sup> field of view. It is found that far-UV to optical collisional lines with critical densities,  $N_{\text{cr}}$ , ranging from  $10^3$  to  $10^9 \text{ cm}^{-3}$  suffer collisional de-excitation near the rest velocity of the proplyd correlating tightly with their critical densities. Lines of low  $N_{\text{cr}}$  are suppressed the most. The bipolar jet arising from LV 2 is spectrally and spatially well-detected in several emission lines. We compute the [O III] electron temperature profile across LV 2 in velocity space and measure steep temperature variations associated with the red-shifted lobe of the jet, possibly being due to a shock discontinuity. From the velocity-resolved analysis the ionized gas near the rest frame of LV 2 has  $T_e = 9200 \pm 800 \text{ K}$  and  $N_e \sim 10^6 \text{ cm}^{-3}$ , while the red-shifted jet lobe has  $T_e \approx 9000 - 10^4 \text{ K}$  and  $N_e \sim 10^6 - 10^7 \text{ cm}^{-3}$ . The jet flow is highly ionized but contains dense semi-neutral clumps emitting neutral oxygen lines. The abundances of  $\text{N}^+$ ,  $\text{O}^{2+}$ ,  $\text{Ne}^{2+}$ ,  $\text{Fe}^{2+}$ ,  $\text{S}^+$ , and  $\text{S}^{2+}$  are measured for the strong red-shifted jet lobe. Iron in the core of LV 2 is depleted by 2.54 dex with respect to solar as a result of sedimentation on dust, whereas the efficient destruction of dust grains in the fast microjet raises its Fe abundance to at least 30 per cent solar. Sulphur does not show evidence of significant depletion on dust, but its abundance both in the core and the jet is only about half solar.

Accepted by Monthly Notices of the Royal Astronomical Society

<http://arxiv.org/pdf/1107.1531v1>

## Ruling out unresolved binaries in five transitional disks

Sílvia Vicente<sup>1</sup>, Bruno Merín<sup>2</sup>, Markus Hartung<sup>3</sup>, Hervé Bouy<sup>4</sup>, Nuria Huélamo<sup>4</sup>, Étienne Artigau<sup>5</sup>, Jean-Charles Augereau<sup>6</sup>, Ewine van Dishoeck<sup>7</sup>, Johan Olofsson<sup>8</sup>, Isa Oliveira<sup>7</sup>, and Timo Prusti<sup>1</sup>

<sup>1</sup> RSSD, European Space Agency (ESTEC), P.O. Box 299, 2200 AG Noordwijk, The Netherlands

<sup>2</sup> Herschel Science Centre, European Space Agency (ESAC), P.O. Box 78, 28691 Villanueva de la Cañada (Madrid), Spain

<sup>3</sup> Gemini Observatory, c/o AURA, Casilla 603, La Serena, Chile

<sup>4</sup> CAB (INTA-CSIC), LAEFF, P.O. Box 78, 28691 Villanueva de la Cañada, Madrid, Spain

<sup>5</sup> Département de Physique and Observatoire du Mont Mégantic, Université de Montréal, C.P. 6128, Succ. Centre-Ville, Montréal, QC H3C 3J7, Canada

<sup>6</sup> Laboratoire d’Astrophysique de Grenoble, Université Joseph Fourier, CNRS, UMR 5571, Grenoble, France

<sup>7</sup> Leiden Observatory, Leiden University, P.O. Box 9513, 2300 RA Leiden, The Netherlands

<sup>8</sup> Max Planck Institute for Astronomy, Königstuhl 17, D-69117 Heidelberg, Germany

E-mail contact: [svicente@rssd.esa.int](mailto:svicente@rssd.esa.int)

*Context.* The presence of unresolved binaries on sub-arcsecond scales could explain the existence of optically thin inner holes or gaps in circumstellar disks, which are commonly referred to as “transitional” or “cold” disks, and it is the first scenario to check before making any other assumptions.

*Aims.* We aim at detecting the presence of companions inside the inner hole/gap region of a sample of five well known transitional disks using spatially-resolved imaging in the near-IR with the VLT/NACO/S13 camera, which probes projected distances from the primary of typically 0.1 to 7 arcsec. The sample includes the stars DoAr 21, HD 135344B (SAO 206462), HR 4796A, T Cha, and TW Hya, spanning ages of less than 1 to 10 Myr, spectral types of A0 to K7,

and hole/gap outer radii of 4 to 100 AU.

*Methods.* In order to enhance the contrast and to avoid saturation at the core of the point-spread function (PSF), we use narrow-band filters at 1.75 and 2.12  $\mu\text{m}$ . The “locally optimized combination of images” (LOCI) algorithm is applied for an optimal speckle noise removal and PSF subtraction, providing an increase of 0.5-1.5 mag in contrast over the classic method.

*Results.* With the proviso that we could have missed companions owing to unfavorable projections, the VLT/NACO observations rule out the presence of unresolved companions down to an inner radius of about  $0''.1$  from the primary in all five transitional disks and with a detection limit of 2 to 5 mag in contrast. In the disk outer regions the detection limits typically reach 8 to 9 mag in contrast and 4.7 mag for T Cha. Hence, the NACO images resolve part of the inner hole/gap region of all disks with the exception of TW Hya, for which the inner hole is only 4 AU. The  $5\sigma$  sensitivity profiles, together with a selected evolutionary model, allow to discard *stellar* companions within the inner hole/gap region of T Cha, and down to the *substellar* regime for HD 135344B and HR 4796A. DoAr 21 is the only object from the sample of five disks for which the NACO images are sensitive enough for a detection of objects less massive than  $\sim 13 M_{\text{Jup}}$  that is, potential giant planets or low-mass brown dwarfs at radii larger than  $\sim 76$  AU ( $0''.63$ ).

*Conclusions.* These new VLT/NACO observations further constrain the origin of the inner opacity cavities to be owing to closer or lower-mass companions or other mechanisms such as giant planet formation, efficient grain growth, and photoevaporation (for DoAr 21 and HR 4796A).

Accepted by Astronomy & Astrophysics

## **Morphologically Complex Protostellar Envelopes: Structure and Kinematics**

**John Tobin**

University of Michigan

500 Church St. 830 Dennison Building, Ann Arbor, MI 48108, USA

Address as of 1 Sept 2011: National Radio Astronomy Observatory, 520 Edgemont Rd., Charlottesville, VA 22903,  
USA

Electronic mail: [jjtobin at umich.edu](mailto:jjtobin@umich.edu)

Ph.D dissertation directed by: Lee Hartmann and Edwin Bergin

Ph.D degree awarded: June 2011

I present an in-depth study of protostars and their surrounding envelopes of dense gas and dust, using a multitude of observational methods to reveal new details of the star formation process. I use mid-infrared imaging from the *Spitzer Space Telescope*, combined with photometry spanning the near-infrared to millimeter wavelengths, to construct a model of the L1527 protostellar system. I modeled both the spectral energy distribution and resolved scattered light images to determine physical properties of the protostellar system. The nature of the apparent central point source in the *Spitzer* images was uncertain until high-resolution L-band imaging from the Gemini observatory resolved the point source into a disk in scattered light, having a radius of 200 AU. Protostellar envelopes are also often found to cast shadows against the 8 micron Galactic background in *Spitzer* imaging, enabling direct probes of envelope structure. The shadow images show that the dense envelopes around twenty-two Class 0 protostars are generally morphologically complex from 0.1 pc scales down to  $\sim 1000$  AU; they are often filamentary, and frequently non-axisymmetric. The observed envelope structure indicates a likely origin in turbulent cloud structure rather than a quasi-static/equilibrium formation. The complex envelope structure also may indicate an increased likelihood of fragmentation during collapse, forming close binaries. To further characterize these envelopes, I have observed them in the dense molecular gas tracers  $\text{N}_2\text{H}^+$  and  $\text{NH}_3$ , both of which closely follow the 8 micron extinction morphology. The magnitude of the velocity gradients and envelope complexity on  $\sim 10000$  AU scales indicates that the velocity structure may reflect large-scale infall in addition to the often assumed rotation. Comparisons with three-dimensional filamentary and symmetric rotating collapse models reinforce the interpretation of velocities reflecting large-scale infall, showing that the structure of the envelope must be considered when interpreting the velocity field. To more definitively probe rotation, the kinematic structure on sub-1000 AU scales must be studied, where rotation will certainly be a more prominent component of the velocity field.

<http://www.astro.lsa.umich.edu/~jjtobin/dissertation.pdf>

# Molecular gas and dust influenced by massive protostars - spectral surveys in the far-infrared and submillimeter

Matthijs H.D. van der Wiel

Kapteyn Astronomical Institute, University of Groningen & SRON Netherlands Institute for Space Research  
PO Box 800, 9700AV Groningen, NL

Address as of 1 Jul 2011: Institute for Space Imaging Science, University of Lethbridge, Lethbridge, AB, T1K 3M4,  
Canada

Electronic mail: matthijs.vanderwiel at uleth.ca

Ph.D dissertation directed by: Marco Spaans & Floris van der Tak

Ph.D degree awarded: May 2011

The interstellar material from which new stars form is crucially influenced by these same protostars: gas and dust particles are excited and heated by the protostar. At the same time, the star formation process depends on the gas reservoir: if gas is too warm or dynamically ‘stirred’, it will not accumulate and contribute to the final mass of the forming star. This issue is specifically important in the formation of massive stars ( $> 8 M_{\odot}$ ), emitting the bulk of the  $> 10^3 L_{\odot}$  at UV wavelengths. This radiation impedes mass accretion and alters the chemical composition of the circumstellar material. In this thesis, I study molecular gas and dust in various Galactic star-forming regions.

In chapter 2 I employ mid-infrared *Spitzer Space Telescope* photometry to study MSXDC G048.65. Protostars are identified in this Infrared Dark Cloud (IRDC), using a pre-defined grid of Spectral Energy Distribution models for protostellar systems. Protostars are found in various evolutionary stages, indicating that star formation is an ongoing process in this cloud. The derived stellar masses range from  $\sim 1$  to  $\sim 8 M_{\odot}$ . Hence, in contrast with the popular view that IRDCs are sites of high-mass star formation, I conclude that this particular cloud hosts only low- to intermediate-mass protostars.

The topic of chapters 3 and 4 is the molecular envelope of the high-mass protostar AFGL2591, for which spectral scans have been conducted in two large spectral survey programs: the JCMT Spectral Legacy Survey (SLS, 330–373 GHz) and Chemical *Herschel* Surveys of Star-forming regions (CHESS, 490–1240 GHz). The SLS spectral imaging data yields 35 spatially resolved molecular line maps, showing substructure in CO,  $N_2H^+$ ,  $H_2CO$ , CS,  $C_2H$ , and  $CH_3OH$ . Radiative transfer modeling, adopting various physical descriptions of the circumstellar envelope, shows that the geometry of the envelope needs to provide low-opacity lines of sight for molecular line radiation to escape. Such a geometry likely includes outflow cavities and clumpy substructure at scales smaller than  $10^4$  AU. Space-based *Herschel* CHESS data reveals high- $J$  lines of linear rotor molecules, CO, HCN, HNC, CS and  $HCO^+$ . From a comparison of the ladder of CO line strengths with a radiative transfer model that includes a cavity geometry, direct UV illumination, shocks, and a self-consistent chemical balance, I conclude that one still probes primarily the quiescent outer envelope, even at energy levels up to 300 K. More distinctive probes of UV photodissociation and shocks are expected to be found in species such as  $HCO^+$  or HCN and in even higher- $J$  lines of CO.

Chapter 5 uses SLS data of the Orion Bar to study chemical stratification in this prototypical photodissociation region (PDR). Emission from the  $C_2H$  radical lies close to the ionization front, and  $H_2CO$ , SO, CO and HCN are seen progressively deeper into the cloud. A simple gas-phase PDR model reproduces the observed layering, with the exceptions of the placement and abundances of SO and  $H_2CO$ . The conclusion is that depletion onto dust grains (of sulfur, for example) and production of  $H_2CO$  through grain surface chemistry should be incorporated in the models.

Chapter 6 uses the CH signature toward the luminous high-mass star-forming core NGC6334I, extracted from *Herschel* CHESS data. A remarkably low CH abundance of  $7 \times 10^{-11}$  is derived for the hot core, a factor of several hundred below that in diffuse clouds. In contrast with other molecular tracers, CH gas in the hot cores is inferred to trace quiescent material, not heated or dynamically influenced by outflow activity. Model representations of the CH spectrum indicates the presence of four foreground clouds of cold, absorbing gas. Column densities and line-of-sight velocities are derived for these clouds, two of which are inferred to be related to the NGC6334 complex.

<http://irs.ub.rug.nl/ppn/333852478>



# Confronting FU Orionis outbursts and pre/transitional disks: protostellar/protoplanetary disk structure and evolution

Zhaohuan Zhu

University of Michigan, Ann Arbor, USA

1022 Dennison Build, 500 Church Str, Ann Arbor, MI, 48109, USA

Address as of 1 Aug 2011: 4 Ivy Lane, Peyton Hall, Princeton University, Princeton, NJ 08544, USA

Electronic mail: zhuzh at umich.com

Ph.D dissertation directed by: Lee W Hartmann

Ph.D degree awarded: July 2011

The mechanisms and efficiencies of mass transport in accretion disks can be best constrained by studying their time-dependent behavior. For the first part of my thesis work, I studied outbursts of rapid accretion in protostellar disks. This was motivated by observations of the FU Orionis objects, which are young stellar objects with a sudden increase of their brightness. I first constructed disk radiative transfer models to compare with *Spitzer* observations, and found that the outburst disk region extends to scales of an AU or more, with the inferred limits on the viscosity parameter  $\alpha \sim 0.02-0.2$ . I further analyzed the multi-wavelength high resolution spectra to show the outbursting disk follows Keplerian rotation. I next studied the thermal structure of the disk which leads us to propose that the outburst is due to the thermal activation of the magnetorotational instability (MRI) at  $\sim$ AU scales by the gravitational instability (GI). We carried out both analytic studies and 2-Dimensional radiationhydrodynamic simulations to study the outbursts and constrain the activation of the MRI. Then I extended this work by constructing simplified 1-dimensional time-dependent simulations with infall from the protostellar envelope to study protostellar disk formation and long term evolution. I found that the outbursting behavior of the protostellar disks during the infall phase alleviates the so called “luminosity problem”, and the layered accretion at the later phase leads to a massive ‘dead zone’ where planet formation is favored.

In the second part of my thesis work, I studied protoplanetary disks with gaps and holes-so called “pretransitional and transitional disks”, - to study planet formation in young disks. I have studied gap opening by dynamically-interacting multiple giant planets with two-dimensional hydrodynamic simulations. I found that even with as many as four giant planets, additional substantial dust depletion (e.g. growth) is required to explain these gaps and holes, which sheds light on the early planet formation environment.

<http://www.astro.lsa.umich.edu/~zhuzh/>

## *Meetings*

### IAU Symposium 287

## Cosmic Masers: From OH to Ho

January 29 - February 3 2012, Stellenbosch, South Africa

website: <http://iau287.ska.ac.za/> email: [iaus287 at hartrao.ac.za](mailto:iaus287@hartrao.ac.za)

#### Scientific Rationale:

Masers have left their mark across large areas of astrophysical research. They are observed in the interstellar medium near star-forming regions, they can probe the envelopes around dying stars and planetary nebulae and they occur in supernova remnants; they are even observed within the solar system in comets and at cosmological distances in galaxies where water masers have been observed at red shifts,  $z > 2$ .

Due to their brightness and compactness, masers give information on the structure and kinematics of a wide variety of sources on extremely small scales. They have therefore been important targets for high resolution observations with radio interferometers, such as MERLIN in the UK, the European VLBI Network (EVN), the Very Large Array, (VLA) and the Very Long Baseline Array (VLBA) in the US, VERA in Japan, and the Australia Telescope Compact Array (ATCA). VLBI observations of e.g. Galactic water masers give information on the dynamics of star formation regions on scales of less than 0.1 - 1 AU at a distance of 1 kpc, as well as accurate distances to these sources.

We are now contemplating the advent of a new golden age in radio astronomy, with the Sub-millimetre Array (SMA), Herschel, the Korean VLBI Network (KVN), dedicated earlier this year and which will target millimeter wavelength masers, the Atacama Large Millimeter Array (ALMA), upgrades of the Plateau de Bure Interferometer (PdBI), VLA, ATCA and MERLIN and further developments of real-time (e)VLBI and the start of construction of the Square Kilometer Array (SKA) pre-cursors, MeerKAT and ASKAP. Studies of molecular masers are prime goals for these new and improved instruments, so maser research is undergoing a rapid growth and continues to provide answers to many important astrophysical questions.

This meeting will be the fourth international symposium on astrophysical masers. Developments in maser research have accelerated over the past years and will continue to do so while new instruments come online. The first results of ALMA are expected towards the end of 2011. The upgraded EVLA and eMERLIN have already started producing science, while the SKA pre-cursors in Australia (ASKAP) and South Africa (MeerKAT) are on the near horizon, planning their early science. The new capabilities of these instruments for maser research will be highlighted by these early results. By bringing together maser researchers, this meeting aims to lead to the development of new observational programmes intended to answer a range of fundamental astrophysical questions. The meeting will also provide a forum for maser observers to interact with theorists working on the physics that must be understood for a full analysis of the wealth of new data.

Specific focus areas for this symposium will include:

- Advances in Maser Theory
- Polarisation and magnetic fields
- Star formation masers
- Stellar masers
- Maser Surveys
- Astrometry using masers
- AGN and Megamasers
- Cosmology and the Hubble Constant
- New masers and other developments in maser physics
- Masers with new mm facilities

Confirmed Review Speakers:

Anna Bartkiewicz, Shuji Deguchi, Christian Henkel, Karl Menten, Mark Reid, Anita Richards, Vladimir Strelitski, Andrea Tarchi, Dinh Van Trung

Scientific Organizing Committee:

Anna Bartkiewicz (Torun), Athol Kembell (Illinois), Huib van Langevelde (JIVE/Leiden), Jim Moran (Harvard/SAO), Jessica Chapman (CSIRO), Kee-Tae Kim (KASI), Liz Humphreys (ESO) co-Chair, Malcolm Gray (Manchester), Mareki Honma (NAOJ), Moshe Elitzur (Kentucky), Roy Booth (SA SKA) Chair, Simon Ellingsen (Tasmania), Valentin Bujarrabal (OAN), Wouter Vlemmings (Chalmers) co-Chair, Yolanda Gomez (UNAM)

Local Organizing Committee:

Johan van der Walt (North-West University), Kim de Boer (SKA SA), Michael Gaylard (Hartebeesthoek RAO), Patricia Whitelock (SAAO & UCT), Simon Fishley (SAAO), Anja Schroeder (SKA SA), Nadeem Oozeer (SKA SA), Sean Passmoor (SKA SA), Rose Hames (SKA SA), Roy Booth (SKA SA) Chair, Sharmila Goedhart (SKA SA) co-Chair

Travel Grants:

A limited amount of travel support will be available.

Important dates/deadlines:

Pre-Registration Opens: July 15

Registration Opens: August 1

Application for IAU Travel Support Deadline: September 16

Abstract Submission Deadline: November 1

(with a late deadline in January for new results)

Early Registration Closes: December 1

(after that a late registration fee will be applied)

## From Atoms to Pebbles: Herschel's View of Star and Planet Formation

### How do stars and planets form and evolve in time?

Since mid-2009, the Herschel Space Observatory provides new light on these long-standing astrophysical questions. With its 3.5 m primary mirror, it delivers observations in the 55–671  $\mu\text{m}$  range with unprecedented sensitivity, spatial and spectral resolution, revolutionising our understanding of the star and planet formation processes.

Approaching the three year anniversary of the launch of Herschel, the French Space Agency (CNES) and the Institut de Planétologie et d'Astrophysique de Grenoble (IPAG), are organising an exceptional Symposium: **From Atoms to Pebbles: Herschel's view of Star and Planet Formation**. This event will be held over 3.5 days, from 20<sup>th</sup> to 23<sup>rd</sup> of March 2012, in Grenoble (the heart of the French Alps), France.

The scientific themes of the symposium will be centred around the physical and chemical evolution of gas and dust in star and planet forming environments and on the impact of the low-and high-mass star formation processes on the surroundings. Specific sessions will be dedicated to :

- The pre-collapse phase,
- The protostellar phase,
- Planet-forming circumstellar disks,
- Debris disks, together with their connection to extra-solar planets.

Please consider your participation at this event and reserve the week from 20 to 23 March 2012 in your agenda. More details and key dates will be reported in the second announcement, as well as on the Symposium web site.

**To receive future announcements, you are welcome to provide your email address by going to:**

<http://www.herschel2012.com/newsletter/newsletter.php>

*Scientific Committee:* P. André (CEA/Saclay), P. Caselli (Leeds), F. Casoli (CNES), E. Caux (IRAP), N. Evans (Austin), T. Henning (MPIA), I. Kamp (Groningen), R. Liseau (Onsala), A. Roberge (NASA Goddard), M. Wyatt (Cambridge), J.-C. Augereau, Chair (IPAG)

*Organizing Committee:* J.-C. Augereau (IPAG), A. Bacmann (IPAG), P. Boucheron (CTA EVENTS), C. Ceccarelli (IPAG), B. Darolles (CNES), A. Faure (IPAG), P. Hily-Blant (IPAG), S. Maret (IPAG), F. Ménard (IPAG), C. Monsan (CNES), C. Pinte (IPAG), M. Rouzé (CNES), M. Soulié (CTA EVENTS), M.-H. Sztefek (IPAG), W.-F. Thi (IPAG)

Meeting Website: <http://www.herschel2012.com/information/>

## First Announcement

# New Quests in Stellar Astrophysics. III. A Panchromatic View of Solar-Like Stars, With and Without Planets

This is the third meeting of a series aimed at stressing the importance of stellar physics to understand the properties of stars and stellar systems at different scales. In the first meeting, in 2001, we reviewed the impact of stellar physics on global properties of stellar aggregates up to cosmological scales. In the second meeting, held in 2007, we focused on the UV interval to review our current knowledge on the properties of nearby and distant evolved stellar populations. In the proposed event for 2012 we want to bring out the attention to the multi-wavelength analyses of solar like (H-fueled) stars and address some of the important recent findings related to exoplanet search. Solar like stars have been subjects of analyses by their own right over the years. Nevertheless, it is undeniable that the study of these stars has been boosted by the discovery of hundreds of exoplanets orbiting them. This conference would be a timely and unique opportunity to collect an up to date view of the outcomes from various on-going surveys aimed at studying normal main sequence stars from the UV to radio frequencies, as well as to review our current knowledge on the correlation of stellar properties with the presence of (proto) planets. This fresh panorama will undoubtedly help on the preparation of future studies, mainly those related to exoplanet search.

Among the main topics to be discussed are:

Solar Analogues

Surveys of Solar-Like Stars at (sub-)millimeter wavelengths

Protoplanetary and Debris Disks: models and observations

The Age of Intermediate-type Main Sequence Stars

Star-Planet Connection: Chemical Abundances, Stellar Activity, and Interactions

Stellar Noise in Exoplanet Searches

Other Suns Hosting other Earths

Habitability Amongst Exoplanets

Scientific Organizing Committee:

Natalie Batalha (San Jose State University, USA), Emanuele Bertone (INAOE, Mexico), Piercarlo Bonifacio (Observatoire du Paris, France), John M. Carpenter (Caltech, USA), Miguel Chavez (INAOE, Mexico), David Hughes (INAOE, Mexico), Lisa Kaltenegger (MPI für Astronomie-Heidelberg, Germany), Susana Lizano (CRyA-UNAM, Mexico), Alejandro Lara (Instituto de Geofísica-UNAM, Mexico), Pablo Mauas (IAFE, Argentina), Rafael Rebolo (Instituto de Astrofísica de Canarias, Spain), Antígona Segura (Instituto de Ciencias Nucleares-UNAM, Mexico), Nuno C. Santos (Centro de Astrofísica, Universidade do Porto, Portugal), Mark Wyatt (IoA, University of Cambridge, UK)

Local Organizing Committee

Fernando Cruz, Victor de La Luz, Raul Mujica, Ana Maria Ramirez, Gerardo Ramos, Olga Vega

If you are interested to receive further information please fill-out the web-form at

<http://www.inaoep.mx/~puerto12/registration.html>.

This conference is sponsored by the Instituto Nacional de Astrofísica, Óptica y Electrónica (INAOE) and the Universidad de Guadalajara (UdeG), Mexico.

## **Physics of the Interstellar and Intergalactic Medium**

**Bruce T. Draine**

This is a graduate textbook, written by one of the leading researchers in the field, on the physical properties of gas and dust in between the stars and the galaxies, the radiation and cosmic rays that this medium interacts with, and the effect of gravitation and magnetic fields. The text is clearly written and well illustrated with close to 200 figures. This textbook is likely to be frequently used in graduate courses on the interstellar and intergalactic medium. The book contains the following chapters:

1. Introduction
2. Collisional Processes
3. Statistical Mechanics and Thermodynamic Equilibrium
4. Energy Levels of Atoms and Ions
5. Energy Levels of Molecules
6. Spontaneous Emission, Stimulated Emission, and Absorption
7. Radiative Transfer
8. HI 21-cm Emission and Absorption
9. Absorption Lines: The Curve of Growth
10. Emission and Absorption by a Thermal Plasma
11. Propagation of Radio Waves through the ISM
12. Interstellar Radiation Fields
13. Ionization Processes
14. Recombination of Ions with Electrons
15. Photoionized Gas
16. Ionization in Predominantly Neutral Regions
17. Collisional Excitation
18. Nebular Diagnostics
19. Radiative Trapping
20. Optical Pumping
21. Interstellar Dust: Observed Properties
22. Scattering and Absorption by Small Particles
23. Composition of Interstellar Dust
24. Temperatures of Interstellar Grains
25. Grain Physics: Charging and Sputtering
26. Grain Dynamics
27. Heating and Cooling of HII Regions
28. The Orion HII Region
29. HI Clouds: Observations
30. HI Clouds: Heating and Cooling
31. Molecular Hydrogen
32. Molecular Clouds: Observations
33. Molecular Clouds: Chemistry and Ionization
34. Physical Processes in Hot Gas
35. Fluid Dynamics
36. Shock Waves
37. Ionization/Dissociation Fronts
38. Stellar Winds
39. Effects of Supernovae on the ISM
40. Cosmic Rays and Gamma Rays

- 41. Gravitational Collapse and Star Formation: Theory
- 42. Star Formation: Observations

A supplementary Solutions Manual is available for this book. It is restricted to teachers using the text in courses. For information on how to obtain a copy, refer to: [http://press.princeton.edu/class\\_use/solutions.html](http://press.princeton.edu/class_use/solutions.html)

Endorsements:

"This is the book that I have been waiting for for twenty years. With exceptional clarity, Draine introduces the underlying physics and brings the basic pieces together to describe the multiphase structure of the interstellar and intergalactic medium. Combined with many useful tables and figures, this book will rapidly become a hit with students and researchers alike. It continues the fine tradition of Princeton professors writing seminal books on this topic."—*Ewine van Dishoeck*, Leiden University

"This book is a comprehensive account of the physical processes that take place in the interstellar medium and that determine its behavior. It is likely to become the bible on the subject."—*Alexander Dalgarno*, Harvard University

Princeton University Press, 2011  
Princeton Series in Astrophysics, ISBN 978-0-691-12213-7  
540 pages, Hardback US\$125.00; Paperback US\$65.00  
<http://press.princeton.edu/titles/9499.html>

The Star Formation Newsletter is a vehicle for fast distribution of information of interest for astronomers working on star formation and molecular clouds. You can submit material for the following sections: *Abstracts of recently accepted papers* (only for papers sent to refereed journals), *Abstracts of recently accepted major reviews* (not standard conference contributions), *Dissertation Abstracts* (presenting abstracts of new Ph.D dissertations), *Meetings* (announcing meetings broadly of interest to the star and planet formation and early solar system community), *New Jobs* (advertising jobs specifically aimed towards persons within the areas of the Newsletter), and *Short Announcements* (where you can inform or request information from the community).

**Latex macros for submitting abstracts and dissertation abstracts (by e-mail to [reipurth@ifa.hawaii.edu](mailto:reipurth@ifa.hawaii.edu)) are appended to each issue of the newsletter. You can also submit via the Newsletter web interface at <http://www2.ifa.hawaii.edu/star-formation/index.cfm>**

The Star Formation Newsletter is available on the World Wide Web at <http://www.ifa.hawaii.edu/users/reipurth/newsletter.htm>.

## Moving ... ??

If you move or your e-mail address changes, please send the editor your new address. If the Newsletter bounces back from an address for three consecutive months, the address is deleted from the mailing list.